

LOCAL COHOMOLOGY OF MULTI-REES ALGEBRAS WITH APPLICATIONS TO JOINT REDUCTIONS AND COMPLETE IDEALS

SHREEDEVI K. MASUTI, TONY J. PUTHENPURAKAL, AND J. K. VERMA

Dedicated to Professor Ngo Viet Trung on his sixtieth birthday

ABSTRACT. Let $(R, \mathfrak{m})$ be a Cohen-Macaulay local ring of dimension d and $\mathbf{I} = (I_1, \dots, I_d)$ be $\mathfrak{m}$ -primary ideals in R . We prove that $\lambda_R([H_{(x_{ii}t_i:1 \leq i \leq d)}^d(\mathcal{R}(\mathcal{F}))]_{\mathbf{n}}) < \infty$, for all $\mathbf{n} \in \mathbb{N}^d$, where $\mathcal{F} = \{\mathcal{F}(\mathbf{n}) : \mathbf{n} \in \mathbb{Z}^d\}$ is an $\mathbf{I}$ -admissible filtration and (x_{ij}) is a strict complete reduction of $\mathcal{F}$ and $\mathcal{R}(\mathcal{F})$ is the multi-Rees algebra of $\mathcal{F}$. As a consequence we prove that the normal joint reduction number of I, J, K is zero in an analytically unramified Cohen-Macaulay local ring of dimension 3 if and only if $\bar{e}_3(IJK) - [\bar{e}_3(IJ) + \bar{e}_3(IK) + \bar{e}_3(JK)] + \bar{e}_3(I) + \bar{e}_3(J) + \bar{e}_3(K) = 0$. This generalizes a theorem of Rees on joint reduction number zero in dimension 2. We apply this theorem to generalize a theorem of M. A. Vitulli in dimension 3.

1. INTRODUCTION

The objective of this paper is to explore finite generation of multigraded components of local cohomology modules of extended multi-Rees algebras of $\mathbb{Z}^d$ -graded filtrations of ideals in a d -dimensional Cohen-Macaulay local ring. As a consequence we obtain a satisfactory generalisation in dimension 3 of an important theorem of D. Rees about $\mathbb{N}^2$ -graded filtrations of complete ideals of joint reduction number zero in two-dimensional analytically unramified Cohen-Macaulay local rings [13]. As a consequence of his theorem, Rees reproved Zariski's theorem about products of complete ideals in two-dimensional regular and psuedo-rational local rings and he also obtained a formula for the normal Hilbert polynomial of a complete ideal in a pseudo-rational local ring. An analogue of Rees' theorem in any dimension is not yet known. This paper presents an analogue in dimension 3.

We shall apply our finiteness theorem to provide necessary and sufficient conditions for the integral closure filtration $\{\overline{I^r J^t K^t}\}$ of three $\mathfrak{m}$ -primary ideals in a 3-dimensional analytically unramified

Key words : strict complete reductions, good complete reductions, good joint reductions, multi-Rees algebra, local cohomology, complete ideal, normal Hilbert polynomial, normal joint reduction number zero.

Cohen-Macaulay local ring to have joint reduction number zero. This, in turn, gives us sufficient conditions for power products of complete monomial ideals to be complete in 3-dimensional polynomial rings. This yields a generalization of a theorem of M. A. Vitulli [15] in dimension 3.

We now describe our main results in detail for which we need to recall several definitions and introduce necessary notation. Let $(R, \mathfrak{m})$ be a d -dimensional Noetherian local ring and let $\mathbf{I} = (I_1, I_2, \dots, I_s)$ for ideals $I_1, I_2, \dots, I_s$ of R . For $\mathbf{n} = (n_1, \dots, n_s) \in \mathbb{Z}^s$, let $\mathbf{I}^{\mathbf{n}} = I_1^{n_1} \dots I_s^{n_s}$. An $\mathbf{I}$ -filtration of ideals is a set of ideals $\mathcal{F} = \{\mathcal{F}(\mathbf{n}) : \mathbf{n} \in \mathbb{Z}^s\}$ such that

- (1) $\mathcal{F}(\mathbf{n}) \subseteq \mathcal{F}(\mathbf{m})$ for all $\mathbf{n}, \mathbf{m} \in \mathbb{Z}^s$ and $\mathbf{n} \geq \mathbf{m}$,
- (2) $\mathbf{I}^{\mathbf{n}} \subseteq \mathcal{F}(\mathbf{n})$ for all $\mathbf{n} \in \mathbb{Z}^s$,
- (3) $\mathcal{F}(\mathbf{n})\mathcal{F}(\mathbf{m}) \subseteq \mathcal{F}(\mathbf{n} + \mathbf{m})$ for all $\mathbf{m}, \mathbf{n} \in \mathbb{Z}^s$.

The principal examples of $\mathbb{Z}^s$ -graded filtrations are the *adic filtration* $\{\mathbf{I}^{\mathbf{n}} = I_1^{n_1} I_2^{n_2} \dots I_s^{n_s}\}$ and the *integral closure filtration* $\{\overline{\mathbf{I}^{\mathbf{n}}} = \overline{I_1^{n_1} I_2^{n_2} \dots I_s^{n_s}}\}$. Recall that the integral closure of an ideal J in a ring R denoted by $\overline{J}$ is the ideal

$$\{r \in R \mid r^n + a_1 r^{n-1} + a_2 r^{n-2} + \dots + a_n = 0 \text{ for some } a_i \in J^i \text{ for } i = 1, 2, \dots, n\}.$$

The ideal J is called *complete* or *integrally closed* if $J = \overline{J}$. For indeterminates $t_1, \dots, t_s$ over R , put $\mathbf{t}^{\mathbf{n}} = t_1^{n_1} \dots t_s^{n_s}$. We say $\mathcal{F}$ is an $\mathbf{I}$ -admissible filtration if the extended Rees algebra $\mathcal{R}'(\mathcal{F}) = \bigoplus_{\mathbf{n} \in \mathbb{Z}^s} \mathcal{F}(\mathbf{n}) \mathbf{t}^{\mathbf{n}}$ is a finite module over the extended Rees algebra $\mathcal{R}'(\mathbf{I}) = \bigoplus_{\mathbf{n} \in \mathbb{Z}^s} \mathbf{I}^{\mathbf{n}} \mathbf{t}^{\mathbf{n}}$. The integral closure of $\mathcal{R}'(\mathbf{I})$ in the ring $R[t_1^{\pm 1}, t_2^{\pm 1}, \dots, t_s^{\pm 1}]$ is the extended Rees algebra $\overline{\mathcal{R}'}(\mathbf{I}) = \overline{\mathcal{R}'(\mathcal{F})}$ where $\mathcal{F}(\mathbf{n}) = \overline{\mathbf{I}^{\mathbf{n}}}$ for all $\mathbf{n} \in \mathbb{Z}^s$.

We write $\mathbf{e}_i$ for the vector in $\mathbb{Z}^s$ which has 1 in the i^{th} position and 0 elsewhere for $i = 1, 2, \dots, s$. The zero vector in $\mathbb{Z}^s$ will be denoted by $\mathbf{0}$. Set $\mathbf{e} = \mathbf{e}_1 + \mathbf{e}_2 + \dots + \mathbf{e}_s$. Now assume that $I_1, I_2, \dots, I_s$ are $\mathfrak{m}$ -primary ideals. An $s \times d$ matrix (x_{ij}) where $x_{ij} \in I_i$ for all $j = 1, \dots, d$ and $i = 1, \dots, s$ is called a *complete reduction* of the $\mathbf{I}$ -filtration $\mathcal{F}$ if

$$\mathcal{F}(\mathbf{n} + \mathbf{e}) = (y_1, \dots, y_d) \mathcal{F}(\mathbf{n})$$

for all large $\mathbf{n} \in \mathbb{N}^s$. Here $y_j = \prod_{i=1}^s x_{ij}$ for $j = 1, \dots, d$. We say that (x_{ij}) is a *good complete reduction* of $\mathcal{F}$ if in addition (x_{ij}) satisfies

$$(y_j : j \in A) \cap \mathcal{F}(\mathbf{n}) = (y_j : j \in A) \mathcal{F}(\mathbf{n} - \mathbf{e}) \text{ for } \mathbf{n} \geq |A| \mathbf{e}$$

for every proper subset $A \subsetneq \{1, \dots, d\}$. Here $|A|$ denotes the cardinality of the set A . If $s = d$ then the elements $x_{ii} = a_i$ for $i = 1, 2, \dots, d$ satisfy the equation

$$\mathcal{F}(\mathbf{n}) = a_1 \mathcal{F}(\mathbf{n} - \mathbf{e}_1) + a_2 \mathcal{F}(\mathbf{n} - \mathbf{e}_2) + \dots + a_d \mathcal{F}(\mathbf{n} - \mathbf{e}_d) \text{ for } \mathbf{n} \gg \mathbf{0}.$$

If the above equation holds for $a_i \in I_i$ for $i = 1, 2, \dots, d$ then we say that $(a_1, a_2, \dots, a_d)$ constitutes a *joint reduction* of $\mathcal{F}$. If the above equation holds for all $\mathbf{n} \geq \mathbf{e}$ then we say that the filtration $\mathcal{F}$ has *joint reduction number zero* with respect to the joint reduction $(a_1, a_2, \dots, a_d)$. If in addition, for each proper subset A of $\{1, 2, \dots, d\}$ and $\mathbf{n} \geq \sum_{i \in A} \mathbf{e}_i$

$$(a_i : i \in A) \cap \mathcal{F}(\mathbf{n}) = \sum_{i \in A} a_i \mathcal{F}(\mathbf{n} - \mathbf{e}_i)$$

then we say that $(a_1, a_2, \dots, a_d)$ is a *good joint reduction* of $\mathcal{F}$. We say that (x_{ij}) is a *strict complete reduction* of $\mathcal{F}$ if for all $\mathbf{n} \geq j\mathbf{e}$ and for $j = 1, 2, \dots, d - 1$ we have

$$(y_1, y_2, \dots, y_j) \cap \mathcal{F}(\mathbf{n}) = (y_1, y_2, \dots, y_j) \mathcal{F}(\mathbf{n} - \mathbf{e}).$$

One of the main observations in this paper is that joint reductions can be studied using local cohomology with support in the ideal $\mathcal{I} = (a_1 t_1, a_2 t_2, \dots, a_d t_d)$ of the extended Rees algebra $\mathcal{R}'(\mathcal{F})$. Our main result in section 2 is the following finiteness theorem:

Theorem 1.1 (Finiteness Theorem). *Let $(R, \mathfrak{m})$ be a Cohen-Macaulay local ring of dimension d . Let $\mathcal{F}$ be an $\mathbf{I}$ -admissible filtration of $\mathfrak{m}$ -primary ideals and $M = \mathcal{R}'(\mathcal{F})$. Then*

- (1) $[H_{\mathcal{I}}^i(M)]_{\mathbf{n}} = 0$ for all $\mathbf{n} \gg \mathbf{0}$ and for all $i \geq 0$.
- (2) If $(y_1, y_2, \dots, y_d)$ is a strict complete reduction of $\mathcal{F}$ then $\lambda_R([H_{\mathcal{I}}^d(M)]_{\mathbf{n}}) < \infty$ for all $\mathbf{n} \geq \mathbf{0}$.

In view of the finiteness theorem, it is natural to ask under what conditions, strict complete reductions and good joint reductions exist. Rees proved their existence for the integral closure filtration $\{\overline{I^r J^s}\}$ of $\mathfrak{m}$ -primary ideals I, J in any 2-dimensional analytically unramified Cohen-Macaulay local ring with infinite residue field. We show their existence in dimension 3 under additional hypothesis. For an $\mathbf{I}$ -admissible filtration $\mathcal{F}$ we set

$$G_i(\mathcal{F}) = \bigoplus_{\mathbf{n} \in \mathbb{N}^s} \frac{\mathcal{F}(\mathbf{n})}{\mathcal{F}(\mathbf{n} + \mathbf{e}_i)} \text{ for all } i = 1, 2, \dots, s.$$

For any $\mathbb{Z}^s$ or $\mathbb{N}^s$ -graded algebra R we write R_{++} for the ideal generated by all elements of degree at least $\mathbf{e}$.

Theorem 1.2. *Let $(R, \mathfrak{m})$ be a three dimensional Cohen-Macaulay local ring with infinite residue field and $\mathbf{I} = (I_1, I_2, I_3)$ for $\mathfrak{m}$ -primary ideals I_1, I_2, I_3 in R . Suppose $\mathcal{F}$ is an $\mathbf{I}$ -admissible filtration. Suppose that $G_i(\mathcal{F})_{++}$ has positive depth and $[H_{G_i(\mathcal{F})_{++}}^1(G_i(\mathcal{F}))]_{\mathbf{n}} = 0$ for all $i = 1, 2, 3$ and all $\mathbf{n} + \mathbf{e}_i \geq 0$. Then there exists a good complete reduction (x_{ij}) of $\mathcal{F}$ such that $(x_{i, \sigma(i)} : i = 1, 2, 3)$ is a good joint reduction of $\mathcal{F}$ and for any proper subset A of $\{1, 2, 3\}$ and $i = 1, 2, 3$*

$$(x_{ij} : j \in A) \cap \mathcal{F}(\mathbf{n}) = (x_{ij} : j \in A)\mathcal{F}(\mathbf{n} - \mathbf{e}_i) \quad \text{for all } \mathbf{n} - \mathbf{e}_i \geq 0.$$

The principal application of good joint reductions is a formula for the length of $[H_{\mathcal{I}}^3(\overline{\mathcal{R}'}(\mathcal{F}))]_{(0,0,0)}$. In order to state the next theorem proved in this paper, we recall the notion of the normal Hilbert function and the normal Hilbert polynomial. For an $\mathfrak{m}$ -primary ideal I in an analytically unramified local ring $(R, \mathfrak{m})$ of dimension d , the *normal Hilbert function* of I is the function $\overline{H}_I(n) = \lambda(R/\overline{I}^n)$. There exists a polynomial $\overline{P}_I(x)$ of degree d such that $\overline{P}_I(n) = \overline{H}_I(n)$ for all large n . We write this polynomial in the form

$$\overline{P}_I(x) = \overline{e}_0(I) \binom{x+d-1}{d} - \overline{e}_1(I) \binom{x+d-2}{d-1} + \cdots + (-1)^d \overline{e}_d(I).$$

The polynomial $\overline{P}_I(x)$ is called the *normal Hilbert polynomial* of I . Let $\overline{\mathcal{R}'} := \overline{\mathcal{R}'}(I, J, K)$.

Theorem 1.3. *Let $(R, \mathfrak{m})$ be a 3-dimensional Cohen-Macaulay analytically unramified local ring. Assume that a good complete reduction of the filtration $\{\overline{I^r J^s K^t}\}$ exists. Let (a, b, c) be a good joint reduction of the filtration $\{\overline{I^r J^s K^t}\}$. Then*

$$\lambda_R([H_{(at_1, bt_2, ct_3)}^3(\overline{\mathcal{R}'})]_{(0,0,0)}) = \overline{e}_3(IJK) - [\overline{e}_3(IJ) + \overline{e}_3(IK) + \overline{e}_3(JK)] + \overline{e}_3(I) + \overline{e}_3(J) + \overline{e}_3(K).$$

As a consequence of this theorem, we are able to generalise Rees' Theorem [13] in dimension three. In dimension three we say that the *normal joint reduction number* of I, J, K is zero with respect to (a, b, c) if for all $r, s, t > 0$

$$\overline{I^r J^s K^t} = a \overline{I^{r-1} J^s K^t} + b \overline{I^r J^{s-1} K^t} + c \overline{I^r J^s K^{t-1}}.$$

Theorem 1.4. *Let the assumptions be as in Theorem 1.3. Then following statements are equivalent:*

- (1) $[H_{(at_1, bt_2, ct_3)}^3(\overline{\mathcal{R}'})]_{(0,0,0)} = 0$,
- (2) *The normal joint reduction number of I, J, K is zero with respect to (a, b, c) ,*
- (3) $\overline{e}_3(IJK) - [\overline{e}_3(IJ) + \overline{e}_3(IK) + \overline{e}_3(JK)] + \overline{e}_3(I) + \overline{e}_3(J) + \overline{e}_3(K) = 0$.

Finally we apply our criterion for joint reduction number zero to obtain a generalisation of a theorem of M. A. Vitulli [15] about normal monomial ideals.

Theorem 1.5. *Let k be a field and $R = k[x, y, z]$, $\mathfrak{m} = (x, y, z)$. Suppose that I, J, K are $\mathfrak{m}$ -primary monomial ideals of R such that $I^r J^s K^t$ is complete for all $r + s + t \leq 2$. Then $I^r J^s K^t$ is complete for all $r, s, t \geq 0$.*

Acknowledgement: We thank Parangama Sarkar for a careful reading of the manuscript.

2. FINITENESS OF MULTIGRADED COMPONENTS OF LOCAL COHOMOLOGY MODULES OF REES ALGEBRAS

In [11], authors have derived a formula for $\lambda_R([H_{(at_1, bt_2)}^2(\overline{\mathcal{R}'}(I, J))]_{(r, s)})$ for all integers $r, s \geq 0$, which shows that $\lambda_R([H_{(at_1, bt_2)}^2(\overline{\mathcal{R}'}(I, J))]_{(r, s)}) < \infty$ in an analytically unramified Cohen-Macaulay local ring of dimension 2 for a good joint reduction (a, b) of $\{\overline{I^r J^s}\}$. Here $\overline{\mathcal{R}'}(I, J)$ is the bigraded Rees algebra of the filtration $\{\overline{I^r J^s}\}$. In general $\lambda_R([H_{(at_1, bt_2)}^2(\mathcal{R}'(\mathcal{F}))]_{(r, s)})$ need not be finite for an (I, J) -admissible filtration $\mathcal{F}$ and for a joint reduction (a, b) of $\mathcal{F}$, even if R is regular. In this section we give an example to illustrate this. However, we show that $\lambda_R([H_{(a_1 t_1, \dots, a_d t_d)}^d(\mathcal{R}'(\mathcal{F}))]_{\mathbf{n}}) < \infty$ for all $\mathbf{n} \in \mathbb{N}^d$ in a Cohen-Macaulay local ring of dimension d , where $a_i = x_{ii}$ for a good complete reduction (x_{ij}) of an $\mathbf{I}$ -admissible filtration $\mathcal{F}$.

2.1. Preliminaries. First we need some lemmas.

Lemma 2.1. *Let $R = \bigoplus_{\mathbf{n} \in \mathbb{Z}^s} R_{\mathbf{n}}$ be a Noetherian $\mathbb{Z}^s$ -graded ring. Suppose $E = \bigoplus_{\mathbf{n} \in \mathbb{Z}^s} E_{\mathbf{n}}$ is a $\mathbb{Z}^s$ -graded R -module. Suppose $E_{\mathbf{n}} = 0$ for all $\mathbf{n} \geq \mathbf{m}$.*

- (1) *Let $f \in R_{\mathbf{d}}$, where $\mathbf{d} \geq \mathbf{0}$. Then for all $i \geq 0$ and all $\mathbf{n} \geq \mathbf{m}$,*

$$[H_{(f)}^i(E)]_{\mathbf{n}} = 0. \quad (1)$$

- (2) *Let $f_1, \dots, f_r$ be homogeneous elements of R of degrees $\mathbf{d}_1, \dots, \mathbf{d}_r \in \mathbb{N}^s$. Then for all $i \geq 0$ and all $\mathbf{n} \geq \mathbf{m}$,*

$$[H_{(f_1, \dots, f_r)}^i(E)]_{\mathbf{n}} = 0.$$

- (3) *Assume $R_{\mathbf{0}} = A$ is a local Noetherian ring with maximal ideal $\mathfrak{m}$. If $\text{Supp } E_{\mathbf{n}} \subseteq \{\mathfrak{m}\}$ for all $\mathbf{n} \in \mathbb{Z}^s$ then $\text{Supp } [H_{(f_1, \dots, f_r)}^i(E)]_{\mathbf{n}} \subseteq \{\mathfrak{m}\}$ for all $\mathbf{n} \in \mathbb{Z}^s$.*

Proof. (1) By [1, Exercise 5.1.14], we have an exact sequence

$$0 \longrightarrow H_{(f)}^0(E) \longrightarrow E \longrightarrow E_f \longrightarrow H_{(f)}^1(E) \longrightarrow 0.$$

Since $E_{\mathbf{n}} = 0$ for all $\mathbf{n} \geq \mathbf{m}$, $[H_{(f)}^0(E)]_{\mathbf{n}} = 0$ for all $\mathbf{n} \geq \mathbf{m}$. To show $[H_{(f)}^1(E)]_{\mathbf{n}} = 0$ for all $\mathbf{n} \geq \mathbf{m}$, consider any $z \in [E_f]_{\mathbf{n}}$. Then $z = \frac{b}{f^r}$ for some $r \in \mathbb{N}$ and $b \in E$. Therefore degree $b = \mathbf{n} + r\mathbf{d} \geq \mathbf{n}$. Hence $[E_f]_{\mathbf{n}} = 0$ for all $\mathbf{n} \geq \mathbf{m}$. This gives $[H_{(f)}^1(E)]_{\mathbf{n}} = 0$ for all $\mathbf{n} \geq \mathbf{m}$.

(2) We use induction on r . For $r = 1$ the assertion follows from the equation (1). Assume $r > 1$ and the result is true for $r - 1$. Let $I = (f_1, \dots, f_{r-1})$. By [1, Exercise 5.1.14], we have the exact sequence

$$\cdots \longrightarrow H_I^{i-1}(E) \longrightarrow H_{I,f_r}^{i-1}(E) \longrightarrow H_{(I,f_r)}^i(E) \longrightarrow H_I^i(E) \longrightarrow H_{I,f_r}^i(E) \longrightarrow \cdots.$$

For all $i \geq 0$, we have an exact sequence

$$0 \longrightarrow H_{(f_r)}^1(H_I^{i-1}(E)) \longrightarrow H_{(I,f_r)}^i(E) \longrightarrow H_{(f_r)}^0(H_I^i(E)) \longrightarrow 0. \quad (2)$$

By induction hypothesis

$$[H_I^j(E)]_{\mathbf{n}} = 0 \text{ for all } \mathbf{n} \geq \mathbf{m} \text{ and all } j \geq 0.$$

Therefore from the equation (1), we obtain

$$[H_{(f_r)}^k(H_I^j(E))]_{\mathbf{n}} = 0 \text{ for all } \mathbf{n} \geq \mathbf{m} \text{ and all } k, j \geq 0.$$

Hence $[H_{(I,f_r)}^i(E)]_{\mathbf{n}} = 0$ for all $\mathbf{n} \geq \mathbf{m}$ and all $i \geq 0$.

(3) Let $\mathfrak{p} \neq \mathbf{m}$ be a prime ideal of A . Note that $R_{\mathbf{0}} \setminus \mathfrak{p}$ has elements of degree $\mathbf{0}$. Since $(E_{\mathbf{n}})_{\mathfrak{p}} = 0$ for all $\mathbf{n} \in \mathbb{Z}^s$, $E_{\mathfrak{p}} = 0$. Hence

$$(H_{(f_1, \dots, f_r)}^i(E))_{\mathfrak{p}} = H_{(f_1, \dots, f_r)A_{\mathfrak{p}}}^i(E_{\mathfrak{p}}) = 0.$$

Therefore $([H_{(f_1, \dots, f_r)}^i(E)]_{\mathbf{n}})_{\mathfrak{p}} = 0$ for all $\mathbf{n} \in \mathbb{Z}^s$ and hence $\text{Supp } [H_{(f_1, \dots, f_r)}^i(E)]_{\mathbf{n}} \subseteq \{\mathbf{m}\}$. □

Corollary 2.2. *Let $R = \bigoplus_{\mathbf{n} \in \mathbb{Z}^s} R_{\mathbf{n}}$ be a Noetherian $\mathbb{Z}^s$ -graded ring and let $M = \bigoplus_{\mathbf{n} \in \mathbb{Z}^s} M_{\mathbf{n}}$ be a finitely generated $\mathbb{Z}^s$ -graded R -module. Let q be a graded ideal of R . Suppose there exists $\mathbf{m} \in \mathbb{Z}^s$ such that for all $i \geq 0$ and all $\mathbf{n} \geq \mathbf{m}$,*

$$[H_q^i(M)]_{\mathbf{n}} = 0.$$

Let $f_1, \dots, f_r$ be homogeneous elements with $\deg f_i \in \mathbb{N}^s$ for all i . Then for $q' = (q, f_1, \dots, f_r)$,

$$[H_{q'}^i(M)]_{\mathbf{n}} = 0 \text{ for all } \mathbf{n} \geq \mathbf{m} \text{ and all } i \geq 0.$$

Furthermore, if $R_0 = A$ is a local Noetherian ring and $\lambda_A([H_q^i(M)]_{\mathbf{n}}) < \infty$ for all $\mathbf{n} \in \mathbb{Z}^s$ and all $i \geq 0$ then

$$\text{Supp}[H_{q'}^i(M)]_{\mathbf{n}} \subseteq \{\mathbf{m}\} \text{ for all } \mathbf{n} \in \mathbb{Z}^s \text{ and all } i \geq 0. \quad (3)$$

Proof. It suffices to show the $r = 1$ case. Let $q' = (q, f)$ where $f = f_1$. From (2), we have the exact sequence

$$0 \longrightarrow H_{(f)}^1(H_q^{i-1}(M)) \longrightarrow H_{q'}^i(M) \longrightarrow H_{(f)}^0(H_q^i(M)) \longrightarrow 0. \quad (4)$$

By Lemma 2.1(1), $[H_{q'}^i(M)]_{\mathbf{n}} = 0$ for all $\mathbf{n} \geq \mathbf{m}$ and all $i \geq 0$. To prove (3), localize the exact sequence (4) at a prime ideal $\mathfrak{p}$ of A , $\mathfrak{p} \neq \mathbf{m}$, and use Lemma 2.1(3) to obtain $H_{q'}^i(M)_{\mathfrak{p}} = 0$ which gives $\text{Supp}[H_{q'}^i(M)]_{\mathbf{n}} \subseteq \{\mathbf{m}\}$ for all $\mathbf{n} \in \mathbb{Z}^s$. $\square$

2.2. Local Cohomology modules of extended Rees algebras of a $\mathbb{Z}^d$ -filtration of ideals.

Let $(R, \mathbf{m})$ be a d -dimensional Cohen-Macaulay local ring with infinite residue field in this subsection. Let $\mathbf{I} = (I_1, \dots, I_d)$ for $\mathbf{m}$ -primary ideals $I_1, \dots, I_d$ of R . Let the $d \times d$ matrix (x_{ij}) where $x_{ij} \in I_i$ for all $j = 1, \dots, d$ and $i = 1, \dots, d$ be a complete reduction of the $\mathbf{I}$ -filtration $\mathcal{F}$. Let $y_j = \prod_{i=1}^d x_{ij}$ for $j = 1, \dots, d$. Let $\mathcal{R}(\mathbf{I}) = \bigoplus_{\mathbf{n} \in \mathbb{N}^d} \mathbf{I}^{\mathbf{n}} \mathbf{t}^{\mathbf{n}}$ be the Rees algebra of the filtration $\{\mathbf{I}^{\mathbf{n}}\}$ and $\mathcal{R}(\mathbf{I})_{++} = \bigoplus_{\mathbf{n} \geq \mathbf{e}} \mathbf{I}^{\mathbf{n}} \mathbf{t}^{\mathbf{n}}$ be an ideal in $\mathcal{R}(\mathbf{I})$. Then $q = (y_1 \mathbf{t}^{\mathbf{e}}, \dots, y_d \mathbf{t}^{\mathbf{e}}) \subseteq \mathcal{R}(\mathbf{I})_{++}$. It is clear that $\sqrt{q} = \sqrt{\mathcal{R}(\mathbf{I})_{++}}$. Hence

$$H_{\mathcal{R}(\mathbf{I})_{++}}^i(\mathcal{R}'(\mathcal{F})) = H_q^i(\mathcal{R}'(\mathcal{F}))$$

for all $i \geq 0$. By an easy argument the following result can be derived from [7, Theorem 5.1]. We set $\mathcal{R}_{++} = \mathcal{R}(\mathbf{I})_{++}$.

Proposition 2.3. *Let $\mathcal{F}$ be an admissible $\mathbf{I}$ -filtration. Let $M = \mathcal{R}'(\mathcal{F})$ and $\mathcal{R}' = \mathcal{R}'(\mathbf{I})$. Then*

- (1) $[H_{\mathcal{R}_{++}}^i(M)]_{\mathbf{n}} = 0$ for all $\mathbf{n} \gg \mathbf{0}$.
- (2) $\lambda_R([H_{\mathcal{R}_{++}}^i(M)]_{\mathbf{n}}) < \infty$ for all $\mathbf{n} \in \mathbb{Z}^d$.

Put $a_i = x_{ii}$ for $i = 1, \dots, d$ and $\mathcal{I} = (a_1 t_1, \dots, a_d t_d)$. Then $\mathcal{R}_{++} \subseteq \sqrt{\mathcal{I}}$ and for all $i \geq 0$

$$H_{(\mathcal{I}, \mathcal{R}_{++})}^i(M) = H_{(\mathcal{I}, q)}^i(M) = H_{\mathcal{I}}^i(M).$$

Corollary 2.4. *Let $\mathcal{F}$ be an $\mathbf{I}$ -admissible filtration and $M = \mathcal{R}'(\mathcal{F})$. Then*

$$[H_{\mathcal{I}}^i(M)]_{\mathbf{n}} = 0 \text{ for all } \mathbf{n} \gg \mathbf{0}.$$

Proof. Follows from Corollary 2.2 and Proposition 2.3. $\square$

Let $\mathcal{R}(\mathcal{F}) = \bigoplus_{\mathbf{n} \in \mathbb{N}^g} \mathcal{F}(\mathbf{n}) \mathbf{t}^{\mathbf{n}}$ be the Rees algebra of $\mathbb{N}^g$ -graded filtration $\mathcal{F}$.

Corollary 2.5. *Let $\mathcal{F}$ be an $\mathbf{I}$ -admissible filtration and $M = \mathcal{R}'(\mathcal{F})$. Put $\theta_j = y_j t_1 \dots t_d$ for $j = 1, \dots, d$, $N_j = M/(\theta_1, \dots, \theta_j)M$. Then*

$$(1) [H_{\mathcal{R}_{++}}^i(N_j)]_{\mathbf{n}} = 0 \text{ for all } \mathbf{n} \gg \mathbf{0}, j = 1, \dots, d \text{ and all } i \geq 0.$$

$$(2) [H_{\mathcal{I}}^i(N_j)]_{\mathbf{n}} = 0 \text{ for all } \mathbf{n} \gg \mathbf{0}, j = 1, \dots, d \text{ and all } i \geq 0.$$

Proof. (1) Let $(N_j)_{\geq \mathbf{0}} = \bigoplus_{\mathbf{n} \geq \mathbf{0}} [N_j]_{\mathbf{n}}$. Since $\mathcal{R}(\mathcal{F})$ is a finite $\mathcal{R}(\mathbf{I})$ -module, $(N_j)_{\geq \mathbf{0}}$ is a finite $\mathcal{R}(\mathbf{I})$ -module for all $j = 1, \dots, d$. Hence, from [7, Theorem 2.3], $[H_{\mathcal{R}_{++}}^i((N_j)_{\geq \mathbf{0}})]_{\mathbf{n}} = 0$ for $\mathbf{n} \gg \mathbf{0}$. Therefore, by [7, Proposition 4.5], $[H_{\mathcal{R}_{++}}^i(N_j)]_{\mathbf{n}} = 0$ for $\mathbf{n} \gg \mathbf{0}$.

(2) Follows from part(1) and Corollary 2.2. $\square$

Remark 2.6. (1) Every good complete reduction of $\mathcal{F}$ is strict good complete reduction of $\mathcal{F}$.

(2) If (x_{ij}) is a good complete reduction of the filtration $\mathcal{F}$ then $(x_{i, \sigma(i)})_{i \in S_d}$ is a good joint reduction for every permutation $\sigma \in S_d$. Here S_d is the symmetric group on d letters.

Lemma 2.7. *Let $(y_1, \dots, y_d)$ be a strict complete reduction of $\mathcal{F}$. Then for $j = 1, \dots, d-1$,*

$$\lambda_R[H_{\mathcal{I}}^{d-j}(N_j)]_{\mathbf{n}} < \infty \text{ for all } \mathbf{n} \geq j\mathbf{e}.$$

Proof. Apply descending induction on j . Suppose $j = d-1$. Consider the exact sequence

$$\begin{array}{ccccccc} 0 & \longrightarrow & B^{(d-1)} & \longrightarrow & N_{d-1}(-\mathbf{e}) & \xrightarrow{\theta_d} & N_{d-1} \longrightarrow N_d \longrightarrow 0 \\ & & & & \searrow & & \nearrow \\ & & & & & K^{(d-1)} & \\ & & & & \nearrow & & \searrow \\ & & 0 & & & & 0 \end{array} \quad (5)$$

Note that

$$[N_{d-1}]_{\mathbf{n}} = \frac{\mathcal{F}(\mathbf{n})}{(y_1, \dots, y_{d-1})\mathcal{F}(\mathbf{n} - \mathbf{e})}.$$

We show that $[H_{\mathcal{I}}^i(B^{(d-1)})]_{\mathbf{n}} = 0$ for all $\mathbf{n} \geq d\mathbf{e}$ and all $i \geq 0$. In view of Lemma 2.1, it is enough to show that $[B^{(d-1)}]_{\mathbf{n}} = 0$ for all $\mathbf{n} \geq d\mathbf{e}$. Let $\bar{p} \in [B^{(d-1)}]_{\mathbf{n}}$ for some $p \in \mathcal{F}(\mathbf{n} - \mathbf{e})$, where $\bar{p}$ denotes the image of p in $N_{d-1}(-\mathbf{e})$. Then $y_d p \in (y_1, \dots, y_{d-1})\mathcal{F}(\mathbf{n} - \mathbf{e})$. Since $y_1, \dots, y_d$ is a regular sequence

$$p \in (y_1, \dots, y_{d-1}) \cap \mathcal{F}(\mathbf{n} - \mathbf{e}) = (y_1, \dots, y_{d-1})\mathcal{F}(\mathbf{n} - 2\mathbf{e}) \text{ for all } \mathbf{n} \geq d\mathbf{e}.$$

Hence $\bar{p} = 0$. Thus $[B^{(d-1)}]_{\mathbf{n}} = 0$ for all $\mathbf{n} \geq d\mathbf{e}$. By the exact sequence (5), we get that the sequence

$$\cdots \longrightarrow [H_{\mathcal{I}}^1(B^{(d-1)})]_{\mathbf{n}} \longrightarrow [H_{\mathcal{I}}^1(N_{d-1})]_{\mathbf{n}-\mathbf{e}} \longrightarrow [H_{\mathcal{I}}^1(K^{(d-1)})]_{\mathbf{n}} \longrightarrow [H_{\mathcal{I}}^2(B^{(d-1)})]_{\mathbf{n}} \longrightarrow \cdots$$

is exact. Hence for $\mathbf{n} \geq d\mathbf{e}$ we have,

$$[H_{\mathcal{I}}^1(N_{d-1})]_{\mathbf{n}-\mathbf{e}} \simeq [H_{\mathcal{I}}^1(K^{(d-1)})]_{\mathbf{n}}. \quad (6)$$

By the exact sequence (5), we also have the exact sequence

$$\cdots \longrightarrow [H_{\mathcal{I}}^0(N_d)]_{\mathbf{n}} \longrightarrow [H_{\mathcal{I}}^1(K^{(d-1)})]_{\mathbf{n}} \longrightarrow [H_{\mathcal{I}}^1(N_{d-1})]_{\mathbf{n}} \longrightarrow \cdots. \quad (7)$$

Fix $(d-1)\mathbf{e} \leq \mathbf{m} \in \mathbb{Z}^d$. By Corollary 2.5, there exist an integer $t_0 \in \mathbb{N}$ such that

$$[H_{\mathcal{I}}^1(N_{d-1})]_{\mathbf{m}+t\mathbf{e}} = 0 \text{ for } t \geq t_0 + 1.$$

Therefore $\lambda_R([H_{\mathcal{I}}^1(N_{d-1})]_{\mathbf{m}+t\mathbf{e}}) < \infty$ for $t \gg 0$. We use decreasing induction on t to show that $\lambda_R([H_{\mathcal{I}}^1(N_{d-1})]_{\mathbf{m}+t\mathbf{e}}) < \infty$ for all $t \geq 0$. Assume the result for $t \geq 1$ and prove it for $t-1$. Since

$$[H_{\mathcal{I}}^0(N_d)]_{\mathbf{n}} \subseteq [H_{\mathcal{R}_{++}}^0(N_d)]_{\mathbf{n}} \subseteq [N_d]_{\mathbf{n}} = \frac{\mathcal{F}(\mathbf{n})}{(y_1, \dots, y_d)\mathcal{F}(\mathbf{n} - \mathbf{e})} \text{ for all } \mathbf{n} \in \mathbb{Z}^d,$$

$\lambda_R([H_{\mathcal{I}}^0(N_d)]_{\mathbf{n}}) < \infty$ for all $\mathbf{n} \in \mathbb{Z}^d$. Hence using induction hypothesis and the exact sequence (7), we obtain

$$\lambda_R([H_{\mathcal{I}}^1(K^{(d-1)})]_{\mathbf{m}+t\mathbf{e}}) < \infty.$$

Since $\mathbf{m} + t\mathbf{e} \geq d\mathbf{e}$, from (6), we get $\lambda_R([H_{\mathcal{I}}^1(N_{d-1})]_{\mathbf{m}+(t-1)\mathbf{e}}) < \infty$. Thus $\lambda_R([H_{\mathcal{I}}^1(N_{d-1})]_{\mathbf{m}+t\mathbf{e}}) < \infty$ for all $t \geq 0$ and hence $\lambda_R([H_{\mathcal{I}}^1(N_{d-1})]_{\mathbf{m}}) < \infty$. Therefore $\lambda_R([H_{\mathcal{I}}^1(N_{d-1})]_{\mathbf{m}})$ has finite length for

all $\mathbf{m} \geq (d-1)\mathbf{e}$.

Assume the result for N_j for $j \geq 2$ and we prove it for N_{j-1} . Consider the exact sequence

$$\begin{array}{ccccccc}
 0 & \longrightarrow & B^{(j-1)} & \longrightarrow & N_{j-1}(-\mathbf{e}) & \xrightarrow{\theta_j} & N_{j-1} \longrightarrow 0 \\
 & & & & \searrow & & \nearrow \\
 & & & & K^{(j-1)} & & \\
 & & \nearrow & & \searrow & & \\
 0 & & & & & & 0
 \end{array} \tag{8}$$

Note that

$$[N_{j-1}]_{\mathbf{n}} = \frac{\mathcal{F}(\mathbf{n})}{(y_1, \dots, y_{j-1})\mathcal{F}(\mathbf{n} - \mathbf{e})}.$$

Hence we have the sequence

$$0 \longrightarrow [B^{(j-1)}]_{\mathbf{n}} \longrightarrow \frac{\mathcal{F}(\mathbf{n} - \mathbf{e})}{\sum_{i=1}^{j-1} y_i \mathcal{F}(\mathbf{n} - 2\mathbf{e})} \xrightarrow{y_j} \frac{\mathcal{F}(\mathbf{n})}{\sum_{i=1}^{j-1} y_i \mathcal{F}(\mathbf{n} - \mathbf{e})} \longrightarrow 0.$$

Let $\bar{p} \in [B^{(j-1)}]_{\mathbf{n}}$ for some $p \in \mathcal{F}(\mathbf{n} - \mathbf{e})$ and $py_j \in \sum_{i=1}^{j-1} y_i \mathcal{F}(\mathbf{n} - \mathbf{e})$. Since $y_1, \dots, y_j$ is a regular sequence, we get

$$p \in (y_1, \dots, y_{j-1}) \cap \mathcal{F}(\mathbf{n} - \mathbf{e}) = (y_1, \dots, y_{j-1})\mathcal{F}(\mathbf{n} - 2\mathbf{e}) \text{ for } \mathbf{n} - \mathbf{e} \geq (j-1)\mathbf{e}, \text{ i.e., for } \mathbf{n} \geq j\mathbf{e}.$$

Therefore for $\mathbf{n} \geq j\mathbf{e}$, $[B^{(j-1)}]_{\mathbf{n}} = 0$ which implies that $[H_{\mathcal{I}}^i(B^{(j-1)})]_{\mathbf{n}} = 0$ for all $\mathbf{n} \geq j\mathbf{e}$ and all $i \geq 0$, by Lemma 2.1. By the exact sequence (8), we get

$$\dots \longrightarrow [H_{\mathcal{I}}^{d-j+1}(B^{(j-1)})]_{\mathbf{n}} \longrightarrow [H_{\mathcal{I}}^{d-j+1}(N_{j-1})]_{\mathbf{n}-\mathbf{e}} \longrightarrow [H_{\mathcal{I}}^{d-j+1}(K^{(j-1)})]_{\mathbf{n}} \longrightarrow [H_{\mathcal{I}}^{d-j+2}(B^{(j-1)})]_{\mathbf{n}} \longrightarrow \dots$$

is exact. Hence for all $\mathbf{n} \geq j\mathbf{e}$,

$$[H_{\mathcal{I}}^{d-j+1}(N_{j-1})]_{\mathbf{n}-\mathbf{e}} \simeq [H_{\mathcal{I}}^{d-j+1}(K^{(j-1)})]_{\mathbf{n}}. \tag{9}$$

By the exact sequence (8), we obtain

$$\dots \longrightarrow [H_{\mathcal{I}}^{d-j}(N_j)]_{\mathbf{n}} \longrightarrow [H_{\mathcal{I}}^{d-j+1}(K^{(j-1)})]_{\mathbf{n}} \longrightarrow [H_{\mathcal{I}}^{d-j+1}(N_{j-1})]_{\mathbf{n}} \longrightarrow \dots \tag{10}$$

is exact. By induction hypothesis $[H_{\mathcal{I}}^{d-j}(N_j)]_{\mathbf{n}}$ has finite length for all $\mathbf{n} \geq j\mathbf{e}$. Fix $(j-1)\mathbf{e} \leq \mathbf{m} \in \mathbb{Z}^d$. We use decreasing induction on t to show that $\lambda_R([H_{\mathcal{I}}^{d-j+1}(N_{j-1})]_{\mathbf{m}+t\mathbf{e}}) < \infty$ for all $t \geq 0$. By Corollary 2.5, the result is true for $t \gg 0$. Assume the result for $t \geq 1$ and prove it for $t-1$. As $\mathbf{m} + t\mathbf{e} \geq j\mathbf{e}$, by the exact sequence (10), $[H_{\mathcal{I}}^{d-j+1}(K^{(j-1)})]_{\mathbf{m}+t\mathbf{e}}$ has finite length. Hence,

from (9), $[H_{\mathcal{I}}^{d-j+1}(N_{j-1})]_{\mathbf{m}+(t-1)\mathbf{e}}$ has finite length. Thus $\lambda_R([H_{\mathcal{I}}^{d-j+1}(N_{j-1})]_{\mathbf{m}+t\mathbf{e}}) < \infty$ for all $t \geq 0$ and hence $\lambda_R([H_{\mathcal{I}}^{d-j+1}(N_{j-1})]_{\mathbf{m}}) < \infty$. Therefore $[H_{\mathcal{I}}^{d-j+1}(N_{j-1})]_{\mathbf{n}}$ has finite length for all $\mathbf{n} \geq (j-1)\mathbf{e}$. $\square$

In the following theorem we prove that for a strict complete reduction $(y_1, \dots, y_d)$ of $\mathcal{F}$, $[H_{\mathcal{I}}^d(\mathcal{R}'(\mathcal{F}))]_{\mathbf{n}}$ has finite length for all $\mathbf{n} \geq \mathbf{0}$.

Theorem 2.8. *Let $(R, \mathbf{m})$ be a d -dimensional Cohen-Macaulay local ring. Suppose $\mathbf{I} = (I_1, \dots, I_d)$ is a set of $\mathbf{m}$ -primary ideals and $\mathcal{F}$ is an admissible $\mathbf{I}$ -filtration. Suppose (x_{ij}) is a strict complete reduction of $\mathcal{F}$ and $\mathcal{I} = (a_1 t_1, \dots, a_d t_d)$ be as before. Let $M = \mathcal{R}'(\mathcal{F})$. Then*

$$\lambda_R([H_{\mathcal{I}}^d(M)]_{\mathbf{n}}) < \infty \text{ for all } \mathbf{n} \geq \mathbf{0}.$$

Proof. By Lemma 2.7, $[H_{\mathcal{I}}^{d-1}(N_1)]_{\mathbf{n}}$ has finite length for all $\mathbf{n} \geq \mathbf{e}$. Fix $\mathbf{m} \in \mathbb{N}^d$. Then $\mathbf{m} + t\mathbf{e} \gg \mathbf{0}$ for all large positive integers t . Hence, by Corollary 2.4, there is an integer $t_0 \in \mathbb{N}$ such that

$$[H_{\mathcal{I}}^d(M)]_{\mathbf{m}+t\mathbf{e}} = 0 \text{ for } t \geq t_0 + 1. \quad (11)$$

We use descending induction on t to show that $\lambda_R([H_{\mathcal{I}}^d(M)]_{\mathbf{m}+t\mathbf{e}}) < \infty$. From (11), $\lambda_R([H_{\mathcal{I}}^d(M)]_{\mathbf{m}+t\mathbf{e}}) < \infty$ for $t \geq t_0 + 1$. Assume the result for $t \geq 1$ and prove it for $t - 1$. The exact sequence

$$0 \longrightarrow M(-\mathbf{e}) \xrightarrow{\theta_1} M \longrightarrow N_1 \longrightarrow 0$$

gives the long exact sequence

$$\cdots \longrightarrow [H_{\mathcal{I}}^i(M)]_{\mathbf{m}+(t-1)\mathbf{e}} \longrightarrow [H_{\mathcal{I}}^i(M)]_{\mathbf{m}+t\mathbf{e}} \longrightarrow [H_{\mathcal{I}}^i(N_1)]_{\mathbf{m}+t\mathbf{e}} \longrightarrow \cdots$$

Note that $[H_{\mathcal{I}}^{d-1}(N_1)]_{\mathbf{m}+t\mathbf{e}}$ has finite length. Therefore using the exact sequence

$$\cdots \longrightarrow [H_{\mathcal{I}}^{d-1}(N_1)]_{\mathbf{m}+t\mathbf{e}} \longrightarrow [H_{\mathcal{I}}^d(M)]_{\mathbf{m}+(t-1)\mathbf{e}} \longrightarrow [H_{\mathcal{I}}^d(M)]_{\mathbf{m}+t\mathbf{e}} \longrightarrow \cdots$$

and induction hypothesis we obtain that $\lambda_R([H_{\mathcal{I}}^d(M)]_{\mathbf{m}+(t-1)\mathbf{e}}) < \infty$. Hence $\lambda_R([H_{\mathcal{I}}^d(M)]_{\mathbf{m}+t\mathbf{e}}) < \infty$ for all $t \geq 0$ and thus $\lambda_R([H_{\mathcal{I}}^d(M)]_{\mathbf{m}}) < \infty$. $\square$

We give an example to show that $[H_{(at_1, bt_2)}^2(\mathcal{R}'(I, J))]_{(0,0)}$ need not have finite length in a Cohen-Macaulay local ring of dimension 2 with infinite residue field and a joint reduction (a, b) of (I, J) , where $\mathcal{R}'(I, J)$ is the bigraded Rees algebra of the filtration $\{I^r J^s\}$.

Example 2.9. [3, Example 4.10] Let $R = k[[x, y]]$, $\mathfrak{m} = (x, y)$, $I = (x^2, y^2)$, $J = \mathfrak{m}^2$. Let $a = x^2, b = y^2$. Then, by [3, Example 4.10], $\lambda_R([H_{(at_1, bt_2)}^2(\mathcal{R}'(I, J))]_{(0,0)})$ is not finite. We claim that there does not exist a strict complete reduction of $\{I^r J^s\}$. Suppose there exists a strict complete reduction (x_{ij}) of $\{I^r J^s\}$. Then, by Theorem 2.8, $\lambda_R([H_{(x_{11}t_1, x_{22}t_2)}^2(\mathcal{R}'(I, J))]_{(0,0)}) < \infty$. By [3, Theorem 4.6], it follows that $\lambda_R([H_{(at_1, bt_2)}^2(\mathcal{R}'(I, J))]_{(0,0)}) < \infty$, a contradiction.

3. EXISTENCE OF GOOD COMPLETE REDUCTIONS AND GOOD JOINT REDUCTIONS

Let $(R, \mathfrak{m})$ be a Noetherian local ring of dimension 3 with infinite residue field and $\mathbf{I} = (I_1, I_2, I_3)$ be a set of $\mathfrak{m}$ -primary ideals in R . In this section we prove existence of good complete reductions of an $\mathbf{I}$ -admissible filtration $\mathcal{F}$ under some additional hypothesis on associated graded rings. The following lemma is an analogue of [12, Lemma 1.2] for filtrations of ideals. Since the proof is similar, we skip it.

Lemma 3.1. [Rees' Lemma] Let $(R, \mathfrak{m})$ be a Noetherian local ring of dimension d with infinite residue field and $I_1, \dots, I_g$ be ideals in R . Let $\mathcal{F} = \{\mathcal{F}(\mathbf{n})\}$ be an $\mathbf{I}$ -admissible filtration. Let S be a finite set of prime ideals of R not containing $I_1 \dots I_g$. Then there exist $x_i \in I_i$ not contained in any of the prime ideals in S and integers r_i such that

$$(x_i) \cap \mathcal{F}(\mathbf{n}) = x_i \mathcal{F}(\mathbf{n} - \mathbf{e}_i) \text{ for all } \mathbf{n} \geq r_i \mathbf{e}_i. \quad (12)$$

We put $u_i = t_i^{-1}$ and $u = u_1 \dots u_g$. The proof of [12, Lemma 1.2] shows the following:

Lemma 3.2. Let $\mathcal{F}$ be an $\mathbf{I}$ -admissible filtration. Let $1 \leq i \leq g$ be fixed and

$$S_i = \{\mathfrak{p} \in \text{Ass}_{\mathcal{R}'(\mathcal{F})}(\mathcal{R}'(\mathcal{F})/u\mathcal{R}'(\mathcal{F})) \mid I_i t_i \not\subseteq \mathfrak{p}\}.$$

Suppose there exists an element $x_i \in I_i$ such that $x_i t_i \notin \mathfrak{p}$ for all $\mathfrak{p} \in S_i$. Then there exists an integer r_i such that

$$(x_i) \cap \mathcal{F}(\mathbf{n}) = x_i \mathcal{F}(\mathbf{n} - \mathbf{e}_i) \text{ for all } \mathbf{n} \geq r_i \mathbf{e}_i.$$

Lemma 3.3. Let $(R, \mathfrak{m})$ be a Noetherian local ring of dimension d with infinite residue field and $I_1, \dots, I_g$ be ideals of positive height. Let $\mathcal{F}$ be an $\mathbf{I}$ -admissible filtration. For fixed j , $1 \leq j \leq g$, suppose there exists a nonzerodivisor $x_j \in I_j \setminus \mathfrak{m}I_j$ such that

$$(x_j) \cap \mathcal{F}(\mathbf{n}) = x_j \mathcal{F}(\mathbf{n} - \mathbf{e}_j) \text{ for } n_j \gg 0 \text{ and for all } n_k \geq 0, k \neq j.$$

Let $H_{\mathcal{R}_{++}}^0(G_i(\mathcal{F})) = 0$ for $i = 1, \dots, g$. Then

$$(x_j) \cap \mathcal{F}(\mathbf{n}) = x_j \mathcal{F}(\mathbf{n} - \mathbf{e}_j) \text{ for all } \mathbf{n} \geq \mathbf{e}_j$$

Proof. Since proof is similar for each j , we prove the result for $j = 1$. Let $a = x_1$. Let a^* denote the image of a in $G_i(\mathcal{F})_{\mathbf{e}_1}$. First we prove that $(0 :_{G_i(\mathcal{F})} a^*) = 0$ for each $i = 1, \dots, g$. It suffices to prove the result for $G_1(\mathcal{F})$. For $r \in \mathcal{F}(\mathbf{n})$ let r^* denote the image of r in $[G_1(\mathcal{F})]_{\mathbf{n}}$. Suppose $z^* a^* = 0$ for some $z \in \mathcal{F}(\mathbf{n})$, $\mathbf{n} \geq \mathbf{0}$. Then for any $y^* \in [G_1(\mathcal{F})]_{\mathbf{m}}$, where $y \in \mathcal{F}(\mathbf{m})$, $y^* z^* a^* = 0$. Hence $yza \in (a) \cap \mathcal{F}(\mathbf{m} + \mathbf{n} + 2\mathbf{e}_1) = a\mathcal{F}(\mathbf{m} + \mathbf{n} + \mathbf{e}_1)$ for $m_1 \gg 0$. Thus $y^* z^* = 0$ if $m_1 \gg 0$. Hence $\mathcal{R}_{++}^N z^* = 0$ for some N . Therefore $z^* \in [H_{\mathcal{R}_{++}}^0(G_1(\mathcal{F}))]_{\mathbf{n}}$. Hence $z^* = 0$.

Let $za \in (a) \cap \mathcal{F}(\mathbf{n})$ for $n_1 > 0$. First we prove that $z \in \mathcal{F}(n_2 \mathbf{e}_2)$. Since $\mathcal{F}$ is an $\mathbf{I}$ -admissible filtration, there exists an integer $h \geq 0$ such that $\mathcal{F}(\mathbf{n}) \subseteq \mathbf{I}^{\mathbf{n}-h\mathbf{e}}$ for all $\mathbf{n} \in \mathbb{Z}^g$. Hence $\cap_{k \geq 0} \mathcal{F}(k\mathbf{e}_2) \subseteq \cap_{k \geq h} I_2^{k-h} = 0$. Thus there exists an integer $l \geq 0$, such that $z \in \mathcal{F}(l\mathbf{e}_2) \setminus \mathcal{F}((l+1)\mathbf{e}_2)$. Suppose $l < n_2$. Then $z^* \in [G_2(\mathcal{F})]_{l\mathbf{e}_2}$ and $z^* a^* = 0$ as $za \in \mathcal{F}(\mathbf{n}) \subseteq \mathcal{F}(\mathbf{e}_1 + (l+1)\mathbf{e}_2)$, a contradiction. Therefore $z \in \mathcal{F}(n_2 \mathbf{e}_2)$.

Similar argument shows that $z \in \mathcal{F}(n_2 \mathbf{e}_2 + \dots + n_g \mathbf{e}_g)$. Since

$$\cap_{k \geq 0} \mathcal{F}(k\mathbf{e}_1 + n_2 \mathbf{e}_2 + \dots + n_g \mathbf{e}_g) \subseteq \cap_{k \geq 0} I_1^{k-h} = 0,$$

there exists an integer $l \geq 0$ such that

$$z \in \mathcal{F}(l\mathbf{e}_1 + n_2 \mathbf{e}_2 + \dots + n_g \mathbf{e}_g) \setminus \mathcal{F}((l+1)\mathbf{e}_1 + n_2 \mathbf{e}_2 + \dots + n_g \mathbf{e}_g).$$

Suppose $l < n_1 - 1$. Then $z^* \in [G_1(\mathcal{F})]_{l\mathbf{e}_1 + n_2 \mathbf{e}_2 + \dots + n_g \mathbf{e}_g}$ and $z^* a^* = 0$, a contradiction. Hence $z \in \mathcal{F}(\mathbf{n} - \mathbf{e}_1)$. $\square$

Theorem 3.4. Let $(R, \mathfrak{m})$ be a Cohen-Macaulay local ring of dimension 3 with infinite residue field k and I_1, I_2, I_3 be $\mathfrak{m}$ -primary ideals in R . Let $\mathcal{F}$ be an (I_1, I_2, I_3) -admissible filtration. Assume that for $i = 1, 2, 3$,

$$H_{\mathcal{R}_{++}}^0(G_i(\mathcal{F})) = 0 \text{ and } [H_{\mathcal{R}_{++}}^1(G_i(\mathcal{F}))]_{\mathbf{n}} = 0 \text{ for } \mathbf{n} + \mathbf{e}_i \geq \mathbf{0}.$$

Then there exists a good complete reduction (M_{ij}) , where $M = \begin{bmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ z_1 & z_2 & z_3 \end{bmatrix}$ such that for each $A \subsetneq \{1, 2, 3\}$

$$(x_i : i \in A) \cap \mathcal{F}(r, s, t) = (x_i : i \in A) \mathcal{F}(r-1, s, t) \text{ for } r > 0 \text{ and all } s, t \geq 0$$

$$(y_i : i \in A) \cap \mathcal{F}(r, s, t) = (y_i : i \in A) \mathcal{F}(r, s-1, t) \text{ for } s > 0 \text{ and all } r, t \geq 0$$

$$(z_i : i \in A) \cap \mathcal{F}(r, s, t) = (z_i : i \in A) \mathcal{F}(r, s, t-1) \text{ for } t > 0 \text{ and all } r, s \geq 0.$$

In particular, $(M_{i, \sigma(i)})$ is a good joint reduction of $\mathcal{F}$ for each permutation $\sigma \in S_3$.

Proof. From Lemma 3.1 and 3.3, there exist nonzerodivisors $x_1 \in I_1 \setminus \mathfrak{m}I_1, y_1 \in I_2 \setminus \mathfrak{m}I_2, z_1 \in I_3 \setminus \mathfrak{m}I_3$ such that

$$(x_1) \cap \mathcal{F}(r, s, t) = x_1 \mathcal{F}(r-1, s, t) \text{ for } r > 0 \text{ and all } s, t \geq 0$$

$$(y_1) \cap \mathcal{F}(r, s, t) = y_1 \mathcal{F}(r, s-1, t) \text{ for } s > 0 \text{ and all } r, t \geq 0$$

$$(z_1) \cap \mathcal{F}(r, s, t) = z_1 \mathcal{F}(r, s, t-1) \text{ for } t > 0 \text{ and all } r, s \geq 0.$$

Let $a = x_1 y_1 z_1$, $S_1 = R/(x_1), S_2 = R/(y_1), S_3 = R/(z_1)$ and $S = R/(a)$. Consider the filtration $\mathcal{G} = \{\mathcal{F}(r, s, t)S\}$. Let

$$A_i = \{\mathfrak{p} \in \text{Ass}_{\mathcal{R}'(\mathcal{G})}(\mathcal{R}'(\mathcal{G})/(u\mathcal{R}'(\mathcal{G}))) \mid I_i S t_i \notin \mathfrak{p}\}$$

$$B_i = \{\mathfrak{p} \in \text{Ass}_{\mathcal{R}'(\mathcal{F})}(\mathcal{R}'(\mathcal{F})/(u\mathcal{R}'(\mathcal{F}))) \mid I_i t_i \notin \mathfrak{p}\} \text{ and}$$

$$C_{i,j} = \{\mathfrak{p} \in \text{Ass}_{\mathcal{R}'(\mathcal{F}S_j)}(\mathcal{R}'(\mathcal{F}S_j)/(u\mathcal{R}'(\mathcal{F}S_j))) \mid I_i S_j t_i \notin \mathfrak{p}\}.$$

Then $\frac{\mathfrak{p} \cap I_1 t_1 + \mathfrak{m} I_1 t_1}{\mathfrak{m} I_1 t_1}$ (respectively $W_j = \frac{\mathfrak{p} \cap I_1 S_j t_1 + \mathfrak{m} I_1 S_j t_1}{\mathfrak{m} I_1 S_j t_1}$ and $W = \frac{\mathfrak{p} \cap I_1 S t_1 + \mathfrak{m} I_1 S t_1}{\mathfrak{m} I_1 S t_1}$) is a proper subspace of $\frac{I_1 t_1}{\mathfrak{m} I_1 t_1} \cong \frac{I_1}{\mathfrak{m} I_1}$ (respectively $\frac{I_1 S_j t_1}{\mathfrak{m} I_1 S_j t_1}$ and $\frac{I_1 S t_1}{\mathfrak{m} I_1 S t_1}$) for $\mathfrak{p} \in B_1$ (respectively $\mathfrak{p} \in C_{1,j}$ and $\mathfrak{p} \in A_1$). Let

$$f_j : \frac{I_1}{\mathfrak{m} I_1} \longrightarrow \frac{I_1 S_j}{\mathfrak{m} I_1 S_j} \text{ and } f : \frac{I_1}{\mathfrak{m} I_1} \longrightarrow \frac{I_1 S}{\mathfrak{m} I_1 S}$$

be natural maps of k -vector spaces. Since f_j and f are surjective, $f_j^{-1}(W_j)$ and $f^{-1}(W)$ are proper subspaces of $\frac{I_1}{\mathfrak{m} I_1}$. Since k is infinite, there exists an element $x_2 \in I_1$ such that $x_2 t_1 \notin \mathfrak{p}$ for any $\mathfrak{p} \in B_1, x_2' t_1 \notin \mathfrak{p}$ for any $\mathfrak{p} \in C_{1,j}, A_1$ and x_2 is S -regular. Here $'$ denotes the image of an element in respective quotients. Similarly, there exist $y_2 \in I_2$ and $z_2 \in I_3$ such that $y_2 t_2 \notin \mathfrak{p}$ for

any $\mathfrak{p} \in B_2, y_2' t_2 \notin \mathfrak{p}$ for any $\mathfrak{p} \in A_2, C_{2,j}, z_2 t_3 \notin \mathfrak{p}$ for any $\mathfrak{p} \in B_3, z_2' t_3 \notin \mathfrak{p}$ for any $\mathfrak{p} \in A_3, C_{3,j}$ and y_2, z_2 are nonzerodivisors in S . Therefore by Lemma 3.2 and 3.3

$$\begin{aligned} (x_2) \cap \mathcal{F}(r, s, t) &= x_2 \mathcal{F}(r-1, s, t) \text{ for } r > 0 \text{ and all } s, t \geq 0 \\ x_2 S \cap \mathcal{G}(r, s, t) &= x_2 \mathcal{G}(r-1, s, t) \text{ for } r \gg 0 \text{ and all } s, t \geq 0 \\ x_2 S_j \cap \mathcal{F}(r, s, t) S_j &= x_2 \mathcal{F}(r-1, s, t) S_j \text{ for } r \gg 0 \text{ and all } s, t \geq 0. \end{aligned}$$

Let y_1^* denote the image of y_1 in $[G_i(\mathcal{F})]_{\mathbf{e}_2}$. Note that y_1^* is $G_i(\mathcal{F})$ -regular for each $i = 1, 2, 3$. Therefore we have an exact sequence

$$0 \longrightarrow G_2(\mathcal{F})(0, -1, 0) \xrightarrow{\mu_{y_1^*}} G_2(\mathcal{F}) \longrightarrow G_2(\mathcal{F})/(y_1^*) \longrightarrow 0.$$

This gives the long exact sequence

$$\cdots \longrightarrow H_{\mathcal{R}_{++}}^0(G_2(\mathcal{F})) \longrightarrow H_{\mathcal{R}_{++}}^0(G_2(\mathcal{F})/(y_1^*)) \longrightarrow H_{\mathcal{R}_{++}}^1(G_2(\mathcal{F})(0, -1, 0)) \longrightarrow \cdots.$$

Therefore $[H_{\mathcal{R}_{++}}^0(G_2(\mathcal{F})/(y_1^*))]_{(r,s,t)} = 0$ for all $r, s, t \geq 0$. Consider the exact sequence

$$0 \longrightarrow L \longrightarrow G_2(\mathcal{F})/(y_1^*) \longrightarrow G_2(\mathcal{F}S_2) \longrightarrow 0.$$

Since $(y_1) \cap \mathcal{F}(r, s, t) = y_1 \mathcal{F}(r, s-1, t)$ for $s > 0$, $[G_2(\mathcal{F})/(y_1^*)]_{(r,s,t)} = [G_2(\mathcal{F}S_2)]_{(r,s,t)}$ for $s > 0$.

Thus $L_{(r,s,t)} = 0$ for $s > 0$ and hence L is $\mathcal{R}_{++}$ -torsion. Hence the sequence

$$0 \longrightarrow L \longrightarrow H_{\mathcal{R}_{++}}^0(G_2(\mathcal{F})/(y_1^*)) \longrightarrow H_{\mathcal{R}_{++}}^0(G_2(\mathcal{F}S_2)) \longrightarrow 0$$

is exact. Thus

$$H_{\mathcal{R}_{++}}^0(G_2(\mathcal{F}S_2)) = 0.$$

Considering an exact sequence

$$0 \longrightarrow G_1(\mathcal{F})(0, -1, 0) \xrightarrow{\mu_{y_1^*}} G_1(\mathcal{F}) \longrightarrow G_1(\mathcal{F})/(y_1^*) \longrightarrow 0.$$

we get the long exact sequence

$$\cdots \longrightarrow H_{\mathcal{R}_{++}}^0(G_1(\mathcal{F})) \longrightarrow H_{\mathcal{R}_{++}}^0(G_1(\mathcal{F})/(y_1^*)) \longrightarrow H_{\mathcal{R}_{++}}^1(G_1(\mathcal{F})(0, -1, 0)) \longrightarrow \cdots.$$

This gives $[H_{\mathcal{R}_{++}}^0(G_1(\mathcal{F})/(y_1^*))]_{(r,s,t)} = 0$ for $s \geq 1$ and $r, t \geq 0$. Consider the exact sequence

$$0 \longrightarrow L' \longrightarrow G_1(\mathcal{F})/(y_1^*) \longrightarrow G_1(\mathcal{F}S_2) \longrightarrow 0.$$

Since $(y_1) \cap \mathcal{F}(r, s, t) = y_1 \mathcal{F}(r, s-1, t)$ for $s > 0$, $[G_1(\mathcal{F})/(y_1^*)]_{(r,s,t)} = [G_1(\mathcal{F}S_2)]_{(r,s,t)}$ for $s > 0$. Thus $L'_{(r,s,t)} = 0$ for $s > 0$ and hence L' is $\mathcal{R}_{++}$ -torsion. Hence the sequence

$$0 \longrightarrow L' \longrightarrow H_{\mathcal{R}_{++}}^0(G_1(\mathcal{F})/(y_1^*)) \longrightarrow H_{\mathcal{R}_{++}}^0(G_1(\mathcal{F}S_2)) \longrightarrow 0$$

is exact. Thus

$$[H_{\mathcal{R}_{++}}^0(G_1(\mathcal{F}S_2))]_{(r,s,t)} = 0 \text{ for } s \geq 1.$$

Similar argument shows that

$$[H_{\mathcal{R}_{++}}^0(G_3(\mathcal{F}S_2))]_{(r,s,t)} = 0 \text{ for } s \geq 1.$$

We prove that for all $r, s \geq 1$,

$$(y_1, x_2) \cap \mathcal{F}(r, s, t) = y_1 \mathcal{F}(r, s-1, t) + x_2 \mathcal{F}(r-1, s, t).$$

First we claim that

$$(0 :_{G_2(\mathcal{F}S_2)} (x'_2)^*) = 0 \text{ and } [0 :_{G_i(\mathcal{F}S_2)} (x'_2)^*]_{(r,s,t)} = 0 \text{ if } s \geq 1 \text{ for } i = 1, 3.$$

Here $'$ denotes the image of an element in S_2 . Suppose $(z')^*(x'_2)^* = 0$ for some $(z')^* \in [G_2(\mathcal{F}S_2)]_{(r,s,t)}$. Then $(y')^*(z')^*(x'_2)^* = 0$ for any $(y')^* \in [G_2(\mathcal{F}S_2)]_{(k,l,m)}$. Hence

$$(y z x_2)' \in \mathcal{F}(r+k+1, s+l+1, t+m)S_2 \text{ and } (y z)' \in \mathcal{F}(r+k, s+l+1, t+m)S_2$$

for $k \gg 0$. Thus $(y')^*(z')^* = 0$ in $G_2(\mathcal{F}S_2)$ for $k \gg 0$. Hence $\mathcal{R}_{++}^N(z')^* = 0$ for some N large. Therefore $(z')^* \in [H_{\mathcal{R}_{++}}^0(G_2(\mathcal{F}S_2))]_{(r,s,t)} = 0$.

Suppose $(z')^*(x'_2)^* = 0$ for some $(z')^* \in [G_1(\mathcal{F}S_2)]_{(r,s,t)}$ and $s \geq 1$. Then $(y')^*(z')^*(x'_2)^* = 0$ for any $(y')^* \in [G_1(\mathcal{F}S_2)]_{(k,l,m)}$. Hence $(y z x_2)' \in \mathcal{F}(r+k+2, s+l, t+m)S_2$. Hence $(y z)' \in \mathcal{F}(r+k+1, s+l, t+m)S_2$ for $k \gg 0$. Thus $(y')^*(z')^* = 0$ in $G_1(\mathcal{F}S_2)$ for $k \gg 0$. Hence $\mathcal{R}_{++}^N(z')^* = 0$ for some N large. Therefore $(z')^* \in [H_{\mathcal{R}_{++}}^0(G_1(\mathcal{F}S_2))]_{(r,s,t)} = 0$. Similar argument shows that $[0 :_{G_3(\mathcal{F}S_2)} (x'_2)^*]_{(r,s,t)} = 0$ for $s \geq 1$.

Let $y_1 u + x_2 v \in \mathcal{F}(r, s, t)$. Since $\cap_{n \geq 0} \mathcal{F}(0, n, 0)S_2 \subseteq \cap_{n \geq 0} I_2^{n-h} S_2 = 0$, there exists an integer $l \geq 0$ such that $v' \in \mathcal{F}(0, l, 0)S_2 \setminus \mathcal{F}(0, l+1, 0)S_2$. Suppose $l < s$. Then $x'_2 v' = (x_2 v)' \in \mathcal{F}(r, s, t)S_2 \subseteq \mathcal{F}(1, l+1, 0)S_2$. Hence $(x'_2)^*(v')^* = 0$ in $G_2(\mathcal{F}S_2)$. Thus $0 \neq (v')^* \in (0 :_{G_2(\mathcal{F}S_2)} (x'_2)^*) = 0$, a contradiction. Therefore $l \geq s$. Hence $v' \in \mathcal{F}(0, s, 0)S_2 = \frac{\mathcal{F}(0,s,0) + (y_1)}{(y_1)}$. Therefore $v = v_1 + y_1 v_2$ for some $v_1 \in \mathcal{F}(0, s, 0)$ and $v_2 \in R$. Thus

$$y_1 u + x_2 v = y_1(u + x_2 v_2) + x_2 v_1 \in \mathcal{F}(r, s, t).$$

Suppose $v'_1 \in \mathcal{F}(l, s, 0)S_2 \setminus \mathcal{F}(l+1, s, 0)S_2$ and $l < r-1$. Since $v'_1 x'_2 \in \mathcal{F}(r, s, t)S_2 \subseteq \mathcal{F}(l+2, s, 0)$, $(v'_1)^*(x'_2)^* = 0$ in $G_1(\mathcal{F}S_2)$. Thus $0 \neq (v'_1)^* \in [0 :_{G_1(\mathcal{F}S_2)} (x'_2)^*]_{(r,s,t)}$ for $s \geq 1$, a contradiction. Therefore $l \geq r-1$. Hence $v'_1 \in \mathcal{F}(r-1, s, 0)S_2$. Let $v_1 = v_3 + y_1 v_4$ for some $v_3 \in \mathcal{F}(r-1, s, 0)$ and $v_4 \in R$. Thus

$$y_1 u + x_2 v = y_1(u + x_2 v_2 + x_2 v_4) + x_2 v_3 \in \mathcal{F}(r, s, t).$$

Suppose $v'_3 \in \mathcal{F}(r-1, s, l)S_2 \setminus \mathcal{F}(r-1, s, l+1)S_2$ for some $l < t$. Since $v'_3 x'_2 \in \mathcal{F}(r, s, t)S_2 \subseteq \mathcal{F}(r, s, l+1)S_2$, $(v'_3)^*(x'_2)^* = 0$ in $G_3(\mathcal{F}S_2)$. Thus $0 \neq (v'_3)^* \in [0 :_{G_3(\mathcal{F}S_2)} (x'_2)^*]_{(r,s,t)}$ and $s \geq 1$, a contradiction. Therefore $l \geq t$. Let $v_3 = v_5 + y_1 v_6$ for some $v_5 \in \mathcal{F}(r-1, s, t)$ and $v_6 \in R$. Then

$$y_1 u + x_2 v = y_1(u + x_2 v_2 + x_2 v_4 + x_2 v_6) + x_2 v_5 \in \mathcal{F}(r, s, t).$$

Let $u_1 = u + x_2 v_2 + x_2 v_4 + x_2 v_6$. Since $x_2 v_5 \in \mathcal{F}(r, s, t)$, $y_1 u_1 \in \mathcal{F}(r, s, t) \cap (y_1) = y_1 \mathcal{F}(r, s-1, t)$. Hence $u_1 \in \mathcal{F}(r, s-1, t)$. Thus

$$y_1 u + x_2 v = y_1 u_1 + x_2 v_5 \in y_1 \mathcal{F}(r, s-1, t) + x_2 \mathcal{F}(r-1, s, t).$$

Similar argument shows that

$$\begin{aligned} (y_1, z_2) \cap \mathcal{F}(r, s, t) &= y_1 \mathcal{F}(r, s-1, t) + z_2 \mathcal{F}(r, s, t-1) \text{ for } s, t \geq 1 \text{ and all } r \geq 0 \text{ and} \\ (y_1, y_2) \cap \mathcal{F}(r, s, t) &= (y_1, y_2) \mathcal{F}(r, s-1, t) \text{ for } s \geq 1 \text{ and all } r, t \geq 0. \end{aligned}$$

Similarly, considering the ring S_1 and S_3 , we get

$$\begin{aligned} H_{\mathcal{R}_{++}}^0(G_1(\mathcal{F}S_1)) &= 0 \text{ and } [H_{\mathcal{R}_{++}}^0(G_i(\mathcal{F}S_1))]_{(r,s,t)} = 0 \text{ for all } r \geq 1 \text{ and } i = 2, 3 \text{ and} \\ H_{\mathcal{R}_{++}}^0(G_3(\mathcal{F}S_3)) &= 0 \text{ and } [H_{\mathcal{R}_{++}}^0(G_i(\mathcal{F}S_3))]_{(r,s,t)} = 0 \text{ for all } t \geq 1 \text{ and } i = 1, 2. \end{aligned}$$

Thus

$$\begin{aligned} (x_1, y_2) \cap \mathcal{F}(r, s, t) &= x_1 \mathcal{F}(r-1, s, t) + y_2 \mathcal{F}(r, s-1, t) \text{ for } r, s \geq 1 \text{ and } t \geq 0 \\ (x_1, z_2) \cap \mathcal{F}(r, s, t) &= x_1 \mathcal{F}(r-1, s, t) + z_2 \mathcal{F}(r, s, t-1) \text{ for } r, t \geq 1 \text{ and } s \geq 0 \\ (x_1, x_2) \cap \mathcal{F}(r, s, t) &= (x_1, x_2) \mathcal{F}(r-1, s, t) \text{ for } r \geq 1 \text{ and all } s, t \geq 0 \\ (z_1, x_2) \cap \mathcal{F}(r, s, t) &= z_1 \mathcal{F}(r, s, t-1) + x_2 \mathcal{F}(r-1, s, t) \text{ for } r, t \geq 1 \text{ and } s \geq 0 \\ (z_1, y_2) \cap \mathcal{F}(r, s, t) &= z_1 \mathcal{F}(r, s, t-1) + y_2 \mathcal{F}(r, s-1, t) \text{ for } s, t \geq 1 \text{ and } r \geq 0 \\ (z_1, z_2) \cap \mathcal{F}(r, s, t) &= (z_1, z_2) \mathcal{F}(r, s, t-1) \text{ for } t \geq 1 \text{ and all } r, s \geq 0. \end{aligned}$$

Let $b = x_2y_2z_2$. We claim that

$$(a, b) \cap \mathcal{F}(r, s, t) = (a, b)\mathcal{F}(r-1, s-1, t-1) \text{ for } r, s, t > 1.$$

First we prove $(a) \cap \mathcal{F}(r, s, t) = a\mathcal{F}(r-1, s-1, t-1)$ for $r, s, t > 0$. Let $au \in \mathcal{F}(r, s, t)$. Then $x_1y_1z_1u \in \mathcal{F}(r, s, t)$. Since $r > 0$, $y_1z_1u \in \mathcal{F}(r-1, s, t)$. Since $s > 0$, $z_1u \in \mathcal{F}(r-1, s-1, t)$. As $t > 0$, $u \in \mathcal{F}(r-1, s-1, t-1)$.

Next we prove that $(a, x_2) \cap \mathcal{F}(r, s, t) = a\mathcal{F}(r-1, s-1, t-1) + x_2\mathcal{F}(r-1, s, t)$ for $r, s, t > 0$. Let $ap + x_2q \in \mathcal{F}(r, s, t)$. Hence $x_1y_1z_1p + x_2q \in (y_1, x_2) \cap \mathcal{F}(r, s, t)$. Let $x_1y_1z_1p + x_2q = y_1p_1 + x_2q_1$ for some $p_1 \in \mathcal{F}(r, s-1, t)$ and $q_1 \in \mathcal{F}(r-1, s, t)$. Thus $y_1(x_1z_1p - p_1) \in (x_2)$. Hence $x_1z_1p - p_1 \in (x_2)$. Therefore $p_1 \in (x_2, z_1) \cap \mathcal{F}(r, s-1, t) = x_2\mathcal{F}(r-1, s-1, t) + z_1\mathcal{F}(r, s-1, t-1)$. Let $p_1 = x_2p_2 + z_1p_3$ for some $p_2 \in \mathcal{F}(r-1, s-1, t)$ and $p_3 \in \mathcal{F}(r, s-1, t-1)$. Then

$$ap + x_2q = y_1p_1 + x_2q_1 = y_1z_1p_3 + x_2(q_1 + y_1p_2).$$

Therefore $y_1z_1(x_1p - p_3) \in (x_2)$. Thus $x_1p - p_3 \in (x_2)$. Hence $p_3 \in (x_1, x_2) \cap \mathcal{F}(r, s-1, t-1)$. Let $p_3 = x_1p_4 + x_2p_5$ for some $p_4, p_5 \in \mathcal{F}(r-1, s-1, t-1)$. Thus

$$ap + x_2q = ap_4 + x_2(q_1 + y_1p_2 + y_1z_1p_5) \in a\mathcal{F}(r-1, s-1, t-1) + x_2\mathcal{F}(r-1, s, t).$$

Similar argument shows that

$$(a, y_2) \cap \mathcal{F}(r, s, t) = a\mathcal{F}(r-1, s-1, t-1) + y_2\mathcal{F}(r, s-1, t) \text{ for } r, s, t > 0 \text{ and}$$

$$(a, z_2) \cap \mathcal{F}(r, s, t) = a\mathcal{F}(r-1, s-1, t-1) + z_2\mathcal{F}(r, s, t-1) \text{ for } r, s, t > 0.$$

Next we prove that $(a, b) \cap \mathcal{F}(r, s, t) = (a, b)\mathcal{F}(r-1, s-1, t-1)$ for $r, s, t > 1$. Let $ap + bq \in \mathcal{F}(r, s, t)$. Then $ap + bq \in (a, x_2) \cap \mathcal{F}(r, s, t) = a\mathcal{F}(r-1, s-1, t-1) + x_2\mathcal{F}(r-1, s, t)$. Let $ap + bq = ap_1 + x_2q_1$ for some $p_1 \in \mathcal{F}(r-1, s-1, t-1)$ and $q_1 \in \mathcal{F}(r-1, s, t)$. Then $x_2(y_2z_2q - q_1) \in (a)$. Hence $y_2z_2q - q_1 \in (a)$. Thus $q_1 \in (a, y_2) \cap \mathcal{F}(r-1, s, t) = a\mathcal{F}(r-2, s-1, t-1) + y_2\mathcal{F}(r-1, s-1, t)$. Let $q_1 = aq_2 + y_2q_3$ for some $q_2 \in \mathcal{F}(r-2, s-1, t-1)$ and $q_3 \in \mathcal{F}(r-1, s-1, t)$. Thus

$$ap + bq = a(p_1 + x_2q_2) + x_2y_2q_3.$$

Hence $x_2y_2(z_2q - q_3) \in (a)$. Thus

$$q_3 \in (a, z_2) \cap \mathcal{F}(r-1, s-1, t) = a\mathcal{F}(r-2, s-2, t-1) + z_2\mathcal{F}(r-1, s-1, t-1).$$

Let $q_3 = aq_4 + z_2q_5$ for some $q_4 \in \mathcal{F}(r-2, s-2, t-1)$ and $q_5 \in \mathcal{F}(r-1, s-1, t-1)$. Hence

$$ap + bq = a(p_1 + x_2q_2 + x_2y_2q_4) + bq_5 \in (a, b)\mathcal{F}(r-1, s-1, t-1).$$

Let $T_1 = R/(x_2), T_2 = R/(y_2), T_3 = R/(z_2)$ and $T = R/(a, b)$. Consider the filtration $\mathcal{H} = \{\mathcal{F}(r, s, t)T\}$. Let

$$\begin{aligned} D_i &= \{\mathfrak{p} \in \text{Ass}_{\mathcal{R}'(\mathcal{H})}(\mathcal{R}'(\mathcal{H})/(u\mathcal{R}'(\mathcal{H}))) \mid I_i T t_i \notin \mathfrak{p}\} \text{ and} \\ E_{i,j} &= \{\mathfrak{p} \in \text{Ass}_{\mathcal{R}'(\mathcal{F}T_j)}(\mathcal{R}'(\mathcal{F}T_j)/(u\mathcal{R}'(\mathcal{F}T_j))) \mid I_i T_j t_i \notin \mathfrak{p}\}. \end{aligned}$$

Then there exist $x_3 \in I_1, y_3 \in I_2$ and $z_3 \in I_3$ such that x_3, y_3, z_3 are nonzerodivisors in T and $x_3t_1 \notin \mathfrak{p}$ for any $\mathfrak{p} \in B_1, x'_3t_1 \notin \mathfrak{p}$ for any $\mathfrak{p} \in C_{1,j}, D_1, E_{1,j}, y_3t_2 \notin \mathfrak{p}$ for any $\mathfrak{p} \in B_2, y'_3t_2 \notin \mathfrak{p}$ for any $\mathfrak{p} \in C_{2,j}, D_2, E_{2,j}$ and $z_3t_3 \notin \mathfrak{p}$ for any $\mathfrak{p} \in B_3, z'_3t_3 \notin \mathfrak{p}$ for any $\mathfrak{p} \in C_{3,j}, D_3, E_{3,j}$. Therefore by Lemma 3.2

$$\begin{aligned} (x_3) \cap \mathcal{F}(r, s, t) &= x_3\mathcal{F}(r-1, s, t) \text{ for } r > 0 \text{ and all } s, t \geq 0 \\ x_3S_j \cap \mathcal{F}(r, s, t)S_j &= x_3\mathcal{F}(r-1, s, t)S_j \text{ for } r \gg 0 \text{ and all } s, t \geq 0 \text{ and} \\ x_3T_j \cap \mathcal{F}(r, s, t)T_j &= x_3\mathcal{F}(r-1, s, t)T_j \text{ for } r \gg 0 \text{ and all } s, t \geq 0 \text{ and} \\ x_3T \cap \mathcal{H}(r, s, t) &= x_3\mathcal{H}(r-1, s, t) \text{ for } r \gg 0 \text{ and all } s, t \geq 0. \end{aligned}$$

Then argument as above shows that for $i = 1, 2$

$$\begin{aligned} (y_i, x_3) \cap \mathcal{F}(r, s, t) &= y_i\mathcal{F}(r, s-1, t) + x_3\mathcal{F}(r-1, s, t) \text{ for } r, s > 0 \text{ and all } t \geq 0 \text{ and} \\ (z_i, x_3) \cap \mathcal{F}(r, s, t) &= z_i\mathcal{F}(r, s, t-1) + x_3\mathcal{F}(r-1, s, t) \text{ for } r, t > 0 \text{ and all } s \geq 0 \\ (x_1, x_3) \cap \mathcal{F}(r, s, t) &= (x_1, x_3)\mathcal{F}(r-1, s, t) \text{ for } r \geq 1 \text{ and all } s, t \geq 0 \\ (x_2, x_3) \cap \mathcal{F}(r, s, t) &= (x_2, x_3)\mathcal{F}(r-1, s, t) \text{ for } r \geq 1 \text{ and all } s, t \geq 0. \end{aligned}$$

Similarly y_3, z_3 satisfies required equations. Let $c = x_3y_3z_3$. Then

$$(a, b, c) \cap \mathcal{F}(r, s, t) = (a, b, c)\mathcal{F}(r-1, s-1, t-1) \text{ for all } r, s, t \gg 0.$$

Since $R/(a, b, c)$ is Artinian, $\mathcal{F}(r, s, t) \subseteq (a, b, c)$ for all r, s, t large. Thus

$$\mathcal{F}(r, s, t) = (a, b, c)\mathcal{F}(r-1, s-1, t-1) \text{ for } r, s, t \gg 0.$$

Hence $M = \begin{bmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ z_1 & z_2 & z_3 \end{bmatrix}$ is a good complete reduction of $\mathcal{F}$ with the desired property. $\square$

In what follows we prove that a good complete reduction of $\mathcal{F}$ exists if $\mathcal{R}(\mathcal{F})$ is Cohen-Macaulay. We prove few lemmas required for this purpose. For a homomorphism $\phi : \mathbb{Z}^r \rightarrow \mathbb{Z}^q$, set

$$M^\phi = \bigoplus_{\mathbf{m} \in \mathbb{Z}^q} \left(\bigoplus_{\phi(\mathbf{n})=\mathbf{m}} M_{\mathbf{n}} \right)$$

for any $\mathbb{Z}^r$ -graded module M defined over $\mathbb{Z}^r$ -graded ring T . Then $H_{\mathfrak{a}}^i(M)^\phi = H_{\mathfrak{a}^\phi}^i(M^\phi)$ for any homogeneous ideal $\mathfrak{a}$ in T . See [5].

Lemma 3.5. *Let T be standard $\mathbb{N}^3$ -graded ring defined over a local ring $(R, \mathfrak{m})$ and M be a finitely generated $\mathbb{N}^3$ -graded T -module. Let $\mathcal{M}$ be the maximal homogeneous ideal of T . Suppose that each $j = 1, 2, 3$ satisfies,*

$$[H_{\mathcal{M}}^i(M)]_{\mathbf{n}} = 0 \text{ for all } i \geq 0 \text{ and } n_j \geq 0.$$

Then

$$[H_{T_{++}}^i(M)]_{\mathbf{n}} = 0 \text{ for all } \mathbf{n} \in \mathbb{N}^3 \text{ and all } i \geq 0.$$

Proof. Let $T_i^+ = \bigoplus_{n_i \geq 1} T_{\mathbf{n}}$ and $T_{i,j}^+ = \bigoplus_{n_i, n_j \geq 1} T_{\mathbf{n}}$. Consider a homomorphism $\phi : \mathbb{Z}^3 \rightarrow \mathbb{Z}$ defined as $\phi(r, s, t) = r$. Then M^ϕ is an $\mathbb{N}$ -graded module defined over the $\mathbb{N}$ -graded ring T^ϕ . Let $B_{(r,s)} = T_{(0,r,s)}$ and $B = \bigoplus_{r,s \geq 0} B_{(r,s)}$. Then $\mathcal{N} = \mathfrak{m} \bigoplus_{r+s \geq 1} B_{(r,s)}$ is the maximal homogeneous ideal of B . Then $T^\phi \otimes_B B_{\mathcal{N}}$ is $\mathbb{N}$ -graded ring defined over a local ring $B_{\mathcal{N}}$. We have $[H_{\mathcal{M}^\phi}^i(M^\phi)]_m = \bigoplus_{s,t \in \mathbb{Z}} [H_{\mathcal{M}}^i(M)]_{(m,s,t)}$. Hence $[H_{\mathcal{M}^\phi}^i(M^\phi)]_m = 0$ for all $m \geq 0$ and $i \geq 0$. Therefore

$$[H_{\mathcal{M}^\phi \otimes_B B_{\mathcal{N}}}^i(M^\phi \otimes_B B_{\mathcal{N}})]_m \cong [H_{\mathcal{M}^\phi}^i(M^\phi)]_m \otimes_B B_{\mathcal{N}} = 0 \text{ for all } m \geq 0 \text{ and } i \geq 0.$$

Taking $\mathfrak{a} = 0$ in [5, Lemma 2.3], we get that

$$[H_{(T_1^+)^\phi}^i(M^\phi)]_m \otimes_B B_{\mathcal{N}} \cong [H_{(T_1^+)^\phi \otimes_B B_{\mathcal{N}}}^i(M^\phi \otimes_B B_{\mathcal{N}})]_m = 0 \text{ for all } m \geq 0 \text{ and } i \geq 0.$$

Since $[H_{(T_1^+)^\phi}^i(M^\phi)]_m$ is bigraded B -module and $([H_{(T_1^+)^\phi}^i(M^\phi)]_m)_{\mathcal{N}} = 0$, $[H_{(T_1^+)^\phi}^i(M^\phi)]_m = 0$ for all $m \geq 0$. Since $[H_{(T_1^+)^\phi}^i(M^\phi)]_m = \bigoplus_{s,t \geq 0} [H_{T_1^+}^i(M)]_{(m,s,t)} = 0$,

$$[H_{T_1^+}^i(M)]_{\mathbf{n}} = 0 \text{ for all } n_1 \geq 0 \text{ and } i \geq 0. \quad (13)$$

Taking $\mathfrak{a} = (\bigoplus_{p \geq 1, q \geq 0} B_{(p,q)}) \otimes_B B_{\mathcal{N}}$ in [5, Lemma 2.3], we get that

$$[H_{T_1^+ + T_2^+}^i(M)]_{\mathbf{n}} = 0 \text{ for all } n_1 \geq 0 \text{ and all } i \geq 0. \quad (14)$$

Similar argument shows that

$$[H_{T_i^+}^j(M)]_{\mathbf{n}} = 0 \text{ for all } n_i \geq 0 \text{ and } j \geq 0. \quad (15)$$

Consider $\phi : \mathbb{Z}^3 \rightarrow \mathbb{Z}$ defined as $\phi(r, s, t) = t$. Let $C = \bigoplus_{r,s \geq 0} T_{(r,s,0)}$ and $\mathcal{P}$ be the maximal homogeneous ideal of C . Then taking $\mathfrak{a} = (\bigoplus_{p,q \geq 1} C_{(p,q)}) \otimes_C C_{\mathcal{P}}$ in [5, Lemma 2.3], we get that

$$[H_{T_{1,2}^+ + T_3^+}^i(M)]_{\mathbf{n}} = 0 \text{ for all } n_3 \geq 0 \text{ and all } i \geq 0. \quad (16)$$

Consider the Mayer-Vietoris sequence of local cohomology modules

$$\cdots \rightarrow H_{T_1^+ + T_2^+}^i(M) \rightarrow H_{T_1^+}^i(M) \oplus H_{T_2^+}^i(M) \rightarrow H_{T_{1,2}^+}^i(M) \rightarrow H_{T_1^+ + T_2^+}^{i+1}(M) \rightarrow \cdots$$

Then, by equations (13), (14) and (15), for all $i \geq 0$,

$$[H_{T_{1,2}^+}^i(M)]_{\mathbf{n}} = 0 \text{ if } n_1 \geq 0 \text{ and } n_2 \geq 0. \quad (17)$$

Again considering the Mayer-Vietoris sequence of local cohomology modules

$$\cdots \rightarrow H_{T_{1,2}^+ + T_3^+}^i(M) \rightarrow H_{T_{1,2}^+}^i(M) \oplus H_{T_3^+}^i(M) \rightarrow H_{T_{++}}^i(M) \rightarrow H_{T_{1,2}^+ + T_3^+}^{i+1}(M) \rightarrow \cdots$$

and using equations (15), (17) and (16), we get that $[H_{T_{++}}^i(M)]_{\mathbf{n}} = 0$ for all $\mathbf{n} \in \mathbb{N}^3$ and $i \geq 0$. $\square$

Suppose T is an $\mathbb{N}$ -graded ring defined over a local ring and $\mathcal{M}$ is the maximal homogeneous ideal of T . For $\mathbb{N}$ -graded T -module M let

$$a(M) := \sup\{k \mid [H_{\mathcal{M}}^{\dim M}(M)]_k \neq 0\}.$$

M. Herrmann, E. Hyry and J. Ribbe proved that for homogeneous ideal I of positive height in a multi-graded ring B , $a(\mathcal{R}(I)) = -1$ [4]. The same proof shows the following:

Lemma 3.6. [4, Lemma 2.1] Let B be a multi-graded ring of dimension d defined over a local ring and let $I \subseteq B$ be a homogeneous ideal of positive height. Let $\mathcal{F} = \{I_n\}$ be an I -admissible filtration of homogeneous ideal in B such that $\mathcal{R}(\mathcal{F})$ is a finite $\mathcal{R}(I)$ -module. Then $a(\mathcal{R}(\mathcal{F})) = -1$.

Proof. Let $\mathfrak{n}$ be the maximal homogeneous ideal of B . Let $\mathcal{M} = \mathfrak{n} \bigoplus_{n>0} I^n t^n$ be a maximal homogeneous ideal in $\mathcal{R}(I)$. Then $[H_{\mathcal{M}}^i(\mathcal{R}(\mathcal{F}))]_n = 0$ if and only if $[H_{\mathcal{M} \otimes_B B_{\mathfrak{n}}}^i(\mathcal{R}(\mathcal{F}) \otimes_B B_{\mathfrak{n}})]_n = 0$. Since $\mathcal{R}(\mathcal{F}) \otimes_B B_{\mathfrak{n}} = \mathcal{R}_{B_{\mathfrak{n}}}(\mathcal{F} B_{\mathfrak{n}})$, we may assume that B is local with maximal ideal $\mathfrak{n}$. Let $\mathcal{R}(\mathcal{F})_+ = \bigoplus_{n>0} I_n t^n$ and $G(\mathcal{F}) = \bigoplus_{n \geq 0} I_n / I_{n+1}$. For an $\mathbb{N}$ -graded module M let $M(n)$ denote the graded module M with $M(n)_k = M_{n+k}$. Then we have the exact sequences

$$0 \longrightarrow \mathcal{R}(\mathcal{F})_+ \longrightarrow \mathcal{R}(\mathcal{F}) \longrightarrow B \rightarrow 0$$

$$0 \longrightarrow \mathcal{R}(\mathcal{F})_+(1) \longrightarrow \mathcal{R}(\mathcal{F}) \longrightarrow G(\mathcal{F}) \longrightarrow 0.$$

Therefore we get the long exact sequence of local cohomology modules

$$\cdots \longrightarrow H_{\mathcal{M}}^i(\mathcal{R}(\mathcal{F})_+) \longrightarrow H_{\mathcal{M}}^i(\mathcal{R}(\mathcal{F})) \longrightarrow H_{\mathcal{M}}^i(B) \longrightarrow \cdots \text{ and}$$

$$\cdots \longrightarrow H_{\mathcal{M}}^i(\mathcal{R}(\mathcal{F})_+(1)) \longrightarrow H_{\mathcal{M}}^i(\mathcal{R}(\mathcal{F})) \longrightarrow H_{\mathcal{M}}^i(G(\mathcal{F})) \longrightarrow \cdots .$$

Note that $H_{\mathcal{M}}^i(B) = H_{\mathfrak{n}}^i(B)$. Hence $[H_{\mathcal{M}}^i(B)]_n = 0$ for $n \neq 0$. This gives the isomorphisms

$$[H_{\mathcal{M}}^{d+1}(\mathcal{R}(\mathcal{F})_+)]_n \longrightarrow [H_{\mathcal{M}}^{d+1}(\mathcal{R}(\mathcal{F}))]_n \text{ for } n \neq 0 \quad (18)$$

and the epimorphisms

$$[H_{\mathcal{M}}^{d+1}(\mathcal{R}(\mathcal{F})_+)]_{n+1} \longrightarrow [H_{\mathcal{M}}^{d+1}(\mathcal{R}(\mathcal{F}))]_n \text{ for } n \in \mathbb{Z}. \quad (19)$$

Since $H_{\mathcal{M}}^{d+1}(\mathcal{R}(\mathcal{F})_+)$ is Artinian, $[H_{\mathcal{M}}^{d+1}(\mathcal{R}(\mathcal{F})_+)]_n = 0$ for $n \gg 0$. Therefore, by equations (18) and (19), $[H_{\mathcal{M}}^{d+1}(\mathcal{R}(\mathcal{F}))]_n = 0$ for $n \geq 0$. If $[H_{\mathcal{M}}^{d+1}(\mathcal{R}(\mathcal{F}))]_{-1} = 0$ then $H_{\mathcal{M}}^{d+1}(\mathcal{R}(\mathcal{F})) = 0$, a contradiction. Thus $a(\mathcal{R}(\mathcal{F})) = -1$. $\square$

For a standard $\mathbb{N}^g$ -graded ring T defined over a local ring and finitely generated $\mathbb{N}^g$ -graded T -module M , let

$$a^i(M) = \sup\{k \in \mathbb{Z} \mid [H_{\mathcal{M}}^{\dim M}(M)]_{\mathbf{n}} \neq 0 \text{ for some } \mathbf{n} \in \mathbb{Z}^g \text{ with } n_i = k\}$$

be a -invariants of M [5]. Here $\mathcal{M}$ is the maximal homogeneous ideal of T . E. Hyry proved that $a^i(\mathcal{R}(\mathbf{I})) = -1$ for all $1 \leq i \leq g$ [5, Lemma 2.7]. In following lemma we prove that $a^i(\mathcal{R}(\mathcal{F})) = -1$ for all $1 \leq i \leq g$.

Lemma 3.7. *Let $(R, \mathfrak{m})$ be a Noetherian local ring and $I_1, \dots, I_g$ be $\mathfrak{m}$ -primary ideals in R . Let $\mathcal{F}$ be an $\mathbf{I} = (I_1, \dots, I_g)$ -admissible filtration. Then $a^i(\mathcal{R}(\mathcal{F})) = -1$ for all $1 \leq i \leq g$.*

Proof. Let $\mathcal{G} = \{\mathcal{F}(\mathbf{n} - n_i \mathbf{e}_i)\}$ be $\mathbb{N}^{g-1}$ -graded $(I_1, \dots, \widehat{I_i}, \dots, I_g)$ -filtration. Then $B = \mathcal{R}(\mathcal{G})$ is $\mathbb{N}^{g-1}$ -graded ring and $I = \bigoplus_{\mathbf{m} \in \mathbb{N}^g} \mathbf{I}^{\mathbf{m} - (m_i - 1)\mathbf{e}_i}$ is homogeneous ideal in B . Also, $\mathcal{H} = \{\mathcal{H}(n) = \bigoplus_{\mathbf{m} \in \mathbb{N}^g, m_i = n} \mathcal{F}(\mathbf{m})\}_{n \in \mathbb{N}}$ is an I -filtration in B and $\mathcal{R}_B(\mathcal{H}) = \mathcal{R}(\mathcal{F})$ is a finite $\mathcal{R}_B(I) = \mathcal{R}(I)$ -module. Therefore by Lemma 3.6, $a(\mathcal{R}_B(\mathcal{H})) = -1$. Consider $\phi : \mathbb{Z}^g \rightarrow \mathbb{Z}$ defined as $\phi(\mathbf{n}) = n_i$. Then $[H_{\mathcal{M}}^{d+1}(\mathcal{R}_B(\mathcal{H}))]_n = [H_{\mathcal{M}^\phi}^{d+1}(\mathcal{R}(\mathcal{F}))^\phi]_n = \bigoplus_{\mathbf{m}, m_i = n} [H_{\mathcal{M}}^i(M)]_{\mathbf{m}}$. Hence $a^i(\mathcal{R}(\mathcal{F})) = -1$. $\square$

Lemma 3.8. *Let S^* be a $\mathbb{Z}^g$ -graded ring and let $S = \bigoplus_{\mathbf{n} \in \mathbb{N}^g} S_{\mathbf{n}} \hookrightarrow S^*$ be an inclusion. Then*

- (1) for $i > 1$, $H_{S_{++}}^i(S) \cong H_{S_{++}}^i(S^*)$.
- (2) We have an exact sequence

$$0 \rightarrow H_{S_{++}}^0(S) \rightarrow H_{S_{++}}^0(S^*) \rightarrow S^*/S \rightarrow H_{S_{++}}^1(S) \rightarrow H_{S_{++}}^1(S^*) \rightarrow 0.$$

Proof. We have an exact sequence

$$0 \rightarrow S \rightarrow S^* \rightarrow S^*/S \rightarrow 0.$$

Since S^*/S is S_{++} -torsion, $H_{S_{++}}^0(S^*/S) = S^*/S$ and $H_{S_{++}}^i(S^*/S) = 0$ for all $i > 0$. Considering the long exact sequence of local cohomology modules result follows. $\square$

Theorem 3.9. *Let $(R, \mathfrak{m})$ be a Cohen-Macaulay local ring of dimension 3 with infinite residue field and I, J, K be $\mathfrak{m}$ -primary ideals in R . Let $\mathcal{F} = \{\mathcal{F}(r, s, t)\}$ be an (I, J, K) -admissible filtration. Assume that $\mathcal{F}(\mathbf{n}) = \mathcal{F}(\mathbf{n} - n_i \mathbf{e}_i)$ for $n_i \leq 0$. Suppose $\mathcal{R}(\mathcal{F})$ is Cohen-Macaulay. Then there exists*

a good complete reduction (M_{ij}) , where $M = \begin{bmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ z_1 & z_2 & z_3 \end{bmatrix}$, such that $(M_{i, \sigma(i)})$ is a good joint reduction of $\mathcal{F}$ for each permutation $\sigma \in S_3$.

Proof. By Lemma 3.7, $a^j(\mathcal{R}(\mathcal{F})) = -1$ for $j = 1, 2, 3$. Since $\mathcal{R}(\mathcal{F})$ is Cohen-Macaulay, for each $j = 1, 2, 3$ we have

$$[H_{\mathcal{M}}^i(\mathcal{R}(\mathcal{F}))]_{\mathbf{n}} = 0 \text{ for all } i \geq 0 \text{ and } n_j \geq 0,$$

where $\mathcal{M}$ is the maximal homogeneous ideal of $\mathcal{R}(I, J, K)$. Therefore, by Lemma 3.5, for all $i \geq 0$

$$[H_{\mathcal{R}_{++}}^i(\mathcal{R}(\mathcal{F}))]_{\mathbf{n}} = 0 \text{ for all } \mathbf{n} \in \mathbb{N}^3.$$

Hence, by Lemma 3.8, for all $i \geq 0$

$$[H_{\mathcal{R}_{++}}^i(\mathcal{R}'(\mathcal{F}))]_{\mathbf{n}} = 0 \text{ for all } \mathbf{n} \in \mathbb{N}^3.$$

Consider the exact sequence

$$0 \longrightarrow \mathcal{R}'(\mathcal{F})(\mathbf{e}_i) \xrightarrow{t_i^{-1}} \mathcal{R}'(\mathcal{F}) \longrightarrow G'_i(\mathcal{F}) \longrightarrow 0,$$

where $G'_i(\mathcal{F}) = \mathcal{R}'(\mathcal{F})/t_i^{-1}\mathcal{R}'(\mathcal{F})$. This gives a long exact sequence of local cohomology modules

$$\cdots \longrightarrow [H_{\mathcal{R}_{++}}^j(\mathcal{R}'(\mathcal{F}))]_{\mathbf{n}+\mathbf{e}_i} \longrightarrow [H_{\mathcal{R}_{++}}^j(\mathcal{R}'(\mathcal{F}))]_{\mathbf{n}} \longrightarrow [H_{\mathcal{R}_{++}}^j(G'_i(\mathcal{F}))]_{\mathbf{n}} \longrightarrow [H_{\mathcal{R}_{++}}^{j+1}(\mathcal{R}'(\mathcal{F}))]_{\mathbf{n}+\mathbf{e}_i} \longrightarrow \cdots.$$

Hence $[H_{\mathcal{R}_{++}}^i(G'_i(\mathcal{F}))]_{\mathbf{n}} = 0$ for all $i \geq 0$ and $\mathbf{n} + \mathbf{e}_i \geq \mathbf{0}$. Since $\mathcal{F}(\mathbf{n}) = \mathcal{F}(\mathbf{n} - n_i \mathbf{e}_i)$ for $n_i \leq 0$, $[G'_i(\mathcal{F})]_{\mathbf{n}} = 0$ for $n_i < 0$. Hence $\left[\frac{G'_i(\mathcal{F})}{G_i(\mathcal{F})}\right]_{\mathbf{n}} = 0$ if $\mathbf{n} \geq \mathbf{0}$ or $n_i < 0$. Therefore, by Lemma 3.8, $[H_{\mathcal{R}_{++}}^i(G_i(\mathcal{F}))]_{\mathbf{n}} = [H_{\mathcal{R}_{++}}^i(G'_i(\mathcal{F}))]_{\mathbf{n}}$ for all $i \geq 0$ if $\mathbf{n} \geq \mathbf{0}$ or $n_i < 0$. Hence $[H_{\mathcal{R}_{++}}^i(G_i(\mathcal{F}))]_{\mathbf{n}} = 0$ for all $i \geq 0$ and $\mathbf{n} + \mathbf{e}_i \geq \mathbf{0}$. Hence result follows from Theorem 3.4. $\square$

Example 3.10. Let $R = k[x, y, z]$, $I = (x, y, z)$ and $J = K = (x^2, y, z)$. Then (x, y, z) forms a good joint reduction of $\{I^r J^s K^t\}$.

4. COMPUTATION OF $\lambda_R \left(\frac{\overline{I^r J^s K^t}}{a^r \overline{J^s K^t} + b^s \overline{I^r K^t} + c^t \overline{I^r J^s}} \right)$

Let $(R, \mathfrak{m})$ be an analytically unramified Cohen-Macaulay local ring of dimension 3 and I, J, K be $\mathfrak{m}$ -primary ideals in R . Let (a, b, c) be a good joint reduction of $\{\overline{I^r J^s K^t}\}$. In this section we express the length

$$\lambda_R \left(\frac{\overline{I^r J^s K^t}}{a^r \overline{J^s K^t} + b^s \overline{I^r K^t} + c^t \overline{I^r J^s}} \right)$$

in terms of the normal Hilbert coefficients. The main tool employed for this calculation is the homology of a multigrades version of the Kirby-Mehran complex [8].

Lemma 4.1. *Let $(R, \mathfrak{m})$ be a Noetherian local ring and I, J be ideals in R . Let $a \in I$, $b \in J$ and (a, b) be a regular sequence.*

(1) *Suppose a satisfies*

$$(a) \cap \overline{I^r J^s K^t} = a \overline{I^{r-1} J^s K^t} \text{ for all } r > 0 \text{ and all } s, t \geq 0. \quad (20)$$

Then for all $r \geq 1$ and $s, t \geq 0$,

$$(a^m) \cap \overline{I^r J^s K^t} = a^m \overline{I^{r-m} J^s K^t} \text{ for } 0 < m \leq r.$$

(2) *Suppose (a, b) satisfies*

$$(a, b) \cap \overline{I^r J^s K^t} = a \overline{I^{r-1} J^s K^t} + b \overline{I^r J^{s-1} K^t} \text{ for all } r, s > 0 \text{ and all } t \geq 0. \quad (21)$$

Then for all $r, s \geq 1$ and $t \geq 0$,

$$(a^m, b^n) \cap \overline{I^r J^s K^t} = a^m \overline{I^{r-m} J^s K^t} + b^n \overline{I^r J^{s-n} K^t} \text{ for } 0 < m \leq r, 0 < n \leq s.$$

Proof. (1) Induct on m . From equation (20), result follows for $m = 1$. Let $m + 1 \leq r$ and

$a^{m+1}u \in \overline{I^r J^s K^t}$. Then $a^{m+1}u \in (a^m) \cap \overline{I^r J^s K^t} = a^m \overline{I^{r-m} J^s K^t}$, by induction hypothesis.

Hence $au \in \overline{I^{r-m} J^s K^t} \cap (a) = \overline{a I^{r-m-1} J^s K^t}$. Thus $u \in \overline{I^{r-m-1} J^s K^t}$.

(2) First we use induction on m to prove that $(a^m, b) \cap \overline{I^r J^s K^t} = a^m \overline{I^{r-m} J^s K^t} + b \overline{I^r J^{s-1} K^t}$

for $0 < m \leq r$. From equation (21), the result follows for $m = 1$. Let $1 < m \leq r$ and

$a^m u + bv \in \overline{I^r J^s K^t}$. Thus $a^m u + bv \in \overline{I^r J^s K^t} \cap (a^{m-1}, b) = a^{m-1} \overline{I^{r-(m-1)} J^s K^t} + b \overline{I^r J^{s-1} K^t}$,

by induction hypothesis. Let $a^m u + bv = a^{m-1} p + bq$ for some $p \in \overline{I^{r-(m-1)} J^s K^t}$ and

$q \in \overline{I^r J^{s-1} K^t}$. Then $a^{m-1}(au - p) \in (b)$. Hence $au - p \in (b)$ which implies that $p \in$

$(a, b) \cap \overline{I^{r-(m-1)} J^s K^t} = a \overline{I^{r-m} J^s K^t} + b \overline{I^{r-(m-1)} J^{s-1} K^t}$. Let $p = ap_1 + bp_2$ for some

$p_1 \in \overline{I^{r-m} J^s K^t}$ and $p_2 \in \overline{I^{r-(m-1)} J^{s-1} K^t}$. Thus $a^m u + bv = a^m p_1 + b(a^{m-1} p_2 + q) \in$

$a^m \overline{I^{r-m} J^s K^t} + b \overline{I^r J^{s-1} K^t}$.

Similar argument shows that $(a, b^n) \cap \overline{I^r J^s K^t} = a \overline{I^{r-1} J^s K^t} + b^n \overline{I^r J^{s-n} K^t}$ for $0 < n \leq s$.

To prove the assertion we use induction on $m+n$. From equation (21), the result follows for

$m = n = 1$. Let $m+n > 2$. We may assume that $n > 1$. Let $a^m u + b^n v \in (a^m, b^n) \cap \overline{I^r J^s K^t}$.

Then $a^m u + b^n v = a^m p + bq$ for some $p \in \overline{I^{r-m} J^s K^t}$ and $q \in \overline{I^r J^{s-1} K^t}$. Hence $b(b^{n-1}v -$

$q) \in (a^m)$ which implies that $b^{n-1}v - q \in (a^m)$. Thus $q \in (a^m, b^{n-1}) \cap \overline{I^r J^{s-1} K^t} =$

$a^m \overline{I^{r-m} J^{s-1} K^t} + b^{n-1} \overline{I^r J^{s-n} K^t}$, by induction. Let $q = a^m q_1 + b^{n-1} q_2$ for some $q_1 \in$

$\overline{I^{r-m} J^{s-1} K^t}$ and $q_2 \in \overline{I^r J^{s-n} K^t}$. Thus $a^m u + b^n v = a^m(p + bq_1) + b^n(q_2) \in a^m \overline{I^{r-m} J^s K^t} +$

$b^n \overline{I^r J^{s-n} K^t}$.

□

We now introduce an $\mathbb{N}^3$ -graded version of the Kirby-Mehran complex associated to the joint reduction (a, b, c) of the filtration $\{\overline{I^r J^s K^t}\}$:

$$C((a, b, c), r, s, t) : 0 \longrightarrow \frac{R}{I^r} \oplus \frac{R}{J^s} \oplus \frac{R}{K^t} \xrightarrow{\phi_1} \frac{R}{I^r J^s} \oplus \frac{R}{I^r K^t} \oplus \frac{R}{J^s K^t} \xrightarrow{\phi_0} \frac{R}{I^r J^s K^t} \longrightarrow 0,$$

where ϕ_0 and ϕ_1 are defined as

$$\phi_1(u', v', w') = ((a^r v + b^s u)', (a^r w - c^t u)', (-b^s w - c^t v)'), \phi_0(u', v', w') = (c^t u + b^s v + a^r w)'.$$

Here $'$ denotes image of an element in respective quotients. Let $H_i((a, b, c), r, s, t)$ denote the i th homology of the complex $C((a, b, c), r, s, t)$.

Proposition 4.2. *Let $(R, \mathfrak{m})$ be an analytically unramified Cohen-Macaulay local ring of dimension 3 with infinite residue field and I, J, K be $\mathfrak{m}$ -primary ideals in R . Let (a, b, c) be a good joint reduction of $\{\overline{I^r J^s K^t}\}$. Then for integers $r, s, t > 0$*

- (1) $H_0((a, b, c), r, s, t) \cong \frac{R}{(a^r, b^s, c^t) + \overline{I^r J^s K^t}}$
- (2) $H_1((a, b, c), r, s, t) \cong \frac{\overline{I^r J^s K^t} \cap (a^r, b^s, c^t)}{a^r \overline{J^s K^t} + b^s \overline{I^r K^t} + c^t \overline{I^r J^s}}$
- (3) $H_2((a, b, c), r, s, t) = 0$.

Proof. (1) Since $\text{im } \phi_0 = \frac{(a^r, b^s, c^t) + \overline{I^r J^s K^t}}{\overline{I^r J^s K^t}}$,

$$H_0((a, b, c), r, s, t) \cong \frac{R}{(a^r, b^s, c^t) + \overline{I^r J^s K^t}}.$$

(2) Consider the commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{im } \phi_1 & \longrightarrow & \frac{R}{\overline{I^r J^s}} \oplus \frac{R}{\overline{I^r K^t}} \oplus \frac{R}{\overline{J^s K^t}} & \xrightarrow{\delta} & \frac{(a^r, b^s, c^t)}{a^r \overline{J^s K^t} + b^s \overline{I^r K^t} + c^t \overline{I^r J^s}} \longrightarrow 0 \\ & & \downarrow & & \psi \downarrow & & \gamma \downarrow \\ 0 & \longrightarrow & \ker \phi_0 & \longrightarrow & \frac{R}{\overline{I^r J^s}} \oplus \frac{R}{\overline{I^r K^t}} \oplus \frac{R}{\overline{J^s K^t}} & \xrightarrow{\phi_0} & \frac{(a^r, b^s, c^t) + \overline{I^r J^s K^t}}{\overline{I^r J^s K^t}} \longrightarrow 0 \end{array}$$

where $\delta(u', v', w') = (c^t u + b^s v + a^r w)'$ and γ is the natural map. To prove exactness of top row it is enough to prove exactness at $\frac{R}{\overline{I^r J^s}} \oplus \frac{R}{\overline{I^r K^t}} \oplus \frac{R}{\overline{J^s K^t}}$. Suppose $\delta(u', v', w') = 0$. Then $c^t u + b^s v + a^r w = c^t x + b^s y + a^r z$ for some $x \in \overline{I^r J^s}, y \in \overline{I^r K^t}$ and $z \in \overline{J^s K^t}$. Hence $u - x = b^s x_1 + a^r x_2$ for some $x_1, x_2 \in R$. Similarly, $v - y = a^r y_1 + c^t y_2$ and $w - z = b^s z_1 + c^t z_2$ for some $y_1, y_2, z_1, z_2 \in R$. Thus

$$c^t(b^s x_1 + a^r x_2) + b^s(a^r y_1 + c^t y_2) + a^r(b^s z_1 + c^t z_2) = 0.$$

Hence $x_1 + y_2 \in (a^r), x_2 + z_2 \in (b^s), y_1 + z_1 \in (c^t)$. Therefore $y'_2 = -x'_1$ in $\frac{R}{\overline{I^r}}$, $z'_1 = -y'_1$ in $\frac{R}{\overline{K^t}}$ and $z'_2 = -x'_2$ in $\frac{R}{\overline{J^s}}$. Hence $\phi_1(x'_1, x'_2, y'_1) = (u', v', w')$. Thus $(u', v', w') \in \text{im } \phi_1$.

Using the snake lemma and the fact that ψ is an isomorphism we obtain,

$$H_1((a, b, c), r, s, t) = \frac{\ker \phi_0}{\text{im } \phi_1} \cong \ker \gamma = \frac{\overline{I^r J^s K^t} \cap (a^r, b^s, c^t)}{a^r \overline{J^s K^t} + b^s \overline{I^r K^t} + c^t \overline{I^r J^s}}.$$

(3) Suppose $\phi_1(u', v', w') = 0$. Thus $((a^r v + b^s u)', (a^r w - c^t u)', (-b^s w - c^t v')) = 0$. Hence $a^r v + b^s u \in \overline{I^r J^s} \cap (a^r, b^s) = a^r \overline{J^s} + b^s \overline{I^r}$, by Lemma 4.1. Let $a^r v + b^s u = a^r p + b^s q$ for some

$p \in \overline{J^s}$ and $q \in \overline{I^r}$. Thus $v - p \in (b^s)$ and $u - q \in (a^r)$. Hence $u \in \overline{I^r}$ and $v \in \overline{J^s}$. Therefore $u' = v' = 0$. Similarly, $w' = 0$.

□

Proposition 4.3. *Let $(R, \mathfrak{m})$ be an analytically unramified Cohen-Macaulay local ring of dimension 3 with infinite residue field and I, J, K be $\mathfrak{m}$ -primary ideals in R . Let (a, b, c) be a good joint reduction of $\{\overline{I^r J^s K^t}\}$. Then for $r, s, t > 0$,*

$$\begin{aligned} & \lambda_R \left(\frac{(a^r, b^s, c^t)}{a^r \overline{J^s K^t} + b^s \overline{I^r K^t} + c^t \overline{I^r J^s}} \right) \\ &= \left[\lambda_R \left(\frac{R}{\overline{I^r J^s}} \right) + \lambda_R \left(\frac{R}{\overline{I^r K^t}} \right) + \lambda_R \left(\frac{R}{\overline{J^s K^t}} \right) \right] - \left[\lambda_R \left(\frac{R}{\overline{I^r}} \right) + \lambda_R \left(\frac{R}{\overline{J^s}} \right) + \lambda_R \left(\frac{R}{\overline{K^t}} \right) \right] \end{aligned}$$

Proof. From the complex $C((a, b, c), r, s, t)$, for $r, s, t > 0$, we have

$$\begin{aligned} & \lambda_R \left(\frac{R}{\overline{I^r J^s K^t}} \right) - \left[\lambda_R \left(\frac{R}{\overline{I^r J^s}} \right) + \lambda_R \left(\frac{R}{\overline{I^r K^t}} \right) + \lambda_R \left(\frac{R}{\overline{J^s K^t}} \right) \right] + \left[\lambda_R \left(\frac{R}{\overline{I^r}} \right) + \lambda_R \left(\frac{R}{\overline{J^s}} \right) + \lambda_R \left(\frac{R}{\overline{K^t}} \right) \right] \\ &= \lambda_R(H_0((a, b, c), r, s, t)) - \lambda_R(H_1((a, b, c), r, s, t)) + \lambda_R(H_2((a, b, c), r, s, t)). \end{aligned}$$

Therefore, by Proposition 4.2,

$$\begin{aligned} & \left[\lambda_R \left(\frac{R}{\overline{I^r J^s}} \right) + \lambda_R \left(\frac{R}{\overline{I^r K^t}} \right) + \lambda_R \left(\frac{R}{\overline{J^s K^t}} \right) \right] - \left[\lambda_R \left(\frac{R}{\overline{I^r}} \right) + \lambda_R \left(\frac{R}{\overline{J^s}} \right) + \lambda_R \left(\frac{R}{\overline{K^t}} \right) \right] \\ &= \lambda_R \left(\frac{R}{\overline{I^r J^s K^t}} \right) - \lambda_R \left(\frac{R}{(a^r, b^s, c^t) + \overline{I^r J^s K^t}} \right) + \lambda_R \left(\frac{\overline{I^r J^s K^t} \cap (a^r, b^s, c^t)}{a^r \overline{J^s K^t} + b^s \overline{I^r K^t} + c^t \overline{I^r J^s}} \right) \\ &= \lambda_R \left(\frac{(a^r, b^s, c^t) + \overline{I^r J^s K^t}}{\overline{I^r J^s K^t}} \right) + \lambda_R \left(\frac{\overline{I^r J^s K^t} \cap (a^r, b^s, c^t)}{a^r \overline{J^s K^t} + b^s \overline{I^r K^t} + c^t \overline{I^r J^s}} \right) \\ &= \lambda_R \left(\frac{(a^r, b^s, c^t)}{\overline{I^r J^s K^t} \cap (a^r, b^s, c^t)} \right) + \lambda_R \left(\frac{\overline{I^r J^s K^t} \cap (a^r, b^s, c^t)}{a^r \overline{J^s K^t} + b^s \overline{I^r K^t} + c^t \overline{I^r J^s}} \right) \\ &= \lambda_R \left(\frac{(a^r, b^s, c^t)}{a^r \overline{J^s K^t} + b^s \overline{I^r K^t} + c^t \overline{I^r J^s}} \right). \end{aligned}$$

□

In an analytically unramified local ring of dimension d there exists a polynomial $\overline{P}_{I,J}(x, y) \in \mathbb{Q}[x, y]$ (respectively $\overline{P}_{I,J,K}(x, y, z) \in \mathbb{Q}[x, y, z]$) of total degree d , called as the *normal Hilbert polynomial* of I, J (respectively *normal Hilbert polynomial* of I, J, K), such that $\overline{P}_{I,J}(r, s) = \lambda(R/\overline{I^r J^s})$

(respectively $\overline{P}_{I,J,K}(r, s, t) = \lambda(R/\overline{I^r J^s K^t})$) for $r, s \gg 0$ (respectively $r, s, t \gg 0$). We write

$$\begin{aligned}\overline{P}_{I,J}(x, y) &= \sum_{i+j \leq d} (-1)^{d-(i+j)} \overline{e}_{(i,j)}(I, J) \binom{x+i-1}{i} \binom{y+j-1}{j} \text{ and} \\ \overline{P}_{I,J,K}(x, y, z) &= \sum_{i+j+k \leq d} (-1)^{d-(i+j+k)} \overline{e}_{(i,j,k)} \binom{x+i-1}{i} \binom{y+j-1}{j} \binom{z+k-1}{k}.\end{aligned}$$

Rees proved that $\overline{e}_{(1,0)}(I, J) = \overline{e}_1(I)$ in an analytically unramified Cohen-Macaulay local ring of dimension 2 [13, Theorem 1.2]. In following lemma we derive a similar relation between coefficients of degree 2 in $\overline{P}_{I,J,K}(x, y, z)$, $\overline{P}_{I,J}(x, y)$ and $\overline{P}_I(x)$ in an analytically unramified Cohen-Macaulay local ring of dimension 3.

Lemma 4.4. *Let $(R, \mathfrak{m})$ be an analytically unramified Cohen-Macaulay local ring of dimension 3 and I, J, K be $\mathfrak{m}$ -primary ideals in R . Then we have*

$\overline{e}_{(2,0,0)} = \overline{e}_1(I)$	$\overline{e}_{(0,2,0)} = \overline{e}_1(J)$	$\overline{e}_{(0,0,2)} = \overline{e}_1(K)$
$\overline{e}_{(1,1,0)} = \overline{e}_{(1,1)}(I, J)$	$\overline{e}_{(0,1,1)} = \overline{e}_{(1,1)}(J, K)$	$\overline{e}_{(1,0,1)} = \overline{e}_{(1,1)}(I, K)$

Proof. Let $t > 0$ be fixed. There exists $c \in K$ such that

$$(c) \cap \overline{I^r J^s K^t} = \overline{c I^r J^s K^{t-1}} \text{ for all } r, s \geq 0 \text{ and } t > 0.$$

See [13]. Then, by Lemma 4.1 (1), $(c^t) \cap \overline{I^r J^s K^t} = \overline{c^t I^r J^s}$ for all $t > 0$ and $r, s \geq 0$. Therefore we have the exact sequence

$$0 \longrightarrow \frac{\overline{I^r J^s}}{\overline{I^r J^s K^t}} \longrightarrow \frac{R}{\overline{I^r J^s K^t}} \xrightarrow{\mu_{c^t}} \frac{R}{\overline{I^r J^s K^t}} \longrightarrow \frac{R}{\overline{I^r J^s K^t} + (c^t)} \longrightarrow 0,$$

Therefore $\lambda_R\left(\frac{\overline{I^r J^s}}{\overline{I^r J^s K^t}}\right) = \lambda_R\left(\frac{R}{\overline{I^r J^s K^t} + (c^t)}\right)$. Let $R' = R/(c^t)$. Consider the filtration $\mathcal{F} = \{\mathcal{F}(r, s) : r, s \in \mathbb{Z}\}$, where $\mathcal{F}(r, s) = \left\{\frac{\overline{I^r J^s K^t} + (c^t)}{(c^t)}\right\}$. Let “ r ” denotes image of an ideal in R' . By [12, Theorem 2.4], $\lambda_R\left(\frac{R}{\overline{I^r J^s K^t} + (c^t)}\right)$ is a polynomial of total degree 2 for $r, s \gg 0$ with

$$\lambda_R\left(\frac{R}{\overline{I^r J^s K^t} + (c^t)}\right) = e(I') \binom{r+1}{2} + \overline{e}_{(1,1)}(I', J') rs + e(J') \binom{s+1}{2} + \text{lower degree terms}.$$

Note that $e(I') = t\overline{e}_{(2,0,1)}$, $e(J') = t\overline{e}_{(0,2,1)}$ and $\overline{e}_{(1,1)}(I', J') = t\overline{e}_{(1,1,1)}$. Therefore for $r, s \gg 0$ and $t > 0$,

$$\lambda_R\left(\frac{R}{\overline{I^r J^s K^t} + (c^t)}\right) = t\overline{e}_{(2,0,1)} \binom{r+1}{2} + t\overline{e}_{(1,1,1)} rs + t\overline{e}_{(0,2,1)} \binom{s+1}{2} + \text{lower degree terms}.$$

We have

$$\lambda_R \left(\frac{R}{\overline{I^r J^s K^t}} \right) = \lambda_R \left(\frac{R}{\overline{I^r J^s}} \right) + \lambda_R \left(\frac{\overline{I^r J^s}}{\overline{I^r J^s K^t}} \right).$$

Fixing $t \gg 0$ and comparing the coefficients of degree 2 in r, s , we get

$$\bar{e}_{(2,0,0)} = \bar{e}_{(2,0)}(I, J), \bar{e}_{(1,1,0)} = \bar{e}_{(1,1)}(I, J), \bar{e}_{(0,2,0)} = \bar{e}_{(0,2)}(I, J).$$

By [10, Theorem 4.3.1], $\bar{e}_{(2,0)}(I, J) = \bar{e}_1(I)$ and $\bar{e}_{(0,2)}(I, J) = \bar{e}_1(J)$. See also [13, Theorem 1.2].

Hence $\bar{e}_{(2,0,0)} = \bar{e}_1(I)$, $\bar{e}_{(0,2,0)} = \bar{e}_1(J)$ and $\bar{e}_{(1,1,0)} = \bar{e}_{(1,1)}(I, J)$.

Similar argument shows the other equalities. $\square$

Theorem 4.5. *Let $(R, \mathfrak{m})$ be an analytically unramified Cohen-Macaulay local ring of dimension 3 with infinite residue field and I, J, K be $\mathfrak{m}$ -primary ideals in R . Let (a, b, c) be a good joint reduction of $\{\overline{I^r J^s K^t}\}$. Then for $r, s, t \gg 0$,*

$$\begin{aligned} \lambda_R \left(\frac{\overline{I^r J^s K^t}}{a^r \overline{J^s K^t} + b^s \overline{I^r K^t} + c^t \overline{I^r J^s}} \right) &= r[\bar{e}_{(1,0)}(I, J) + \bar{e}_{(1,0)}(I, K) - \bar{e}_{(1,0,0)} - \bar{e}_2(I)] \\ &+ s[\bar{e}_{(0,1)}(I, J) + \bar{e}_{(0,1)}(J, K) - \bar{e}_{(0,1,0)} - \bar{e}_2(J)] \\ &+ t[\bar{e}_{(0,1)}(I, K) + \bar{e}_{(0,1)}(J, K) - \bar{e}_{(0,0,1)} - \bar{e}_2(K)] \\ &+ \bar{e}_3(IJK) - [\bar{e}_3(IJ) + \bar{e}_3(IK) + \bar{e}_3(JK)] + \bar{e}_3(I) + \bar{e}_3(J) + \bar{e}_3(K). \end{aligned}$$

Proof. We have

$$\begin{aligned} \lambda_R \left(\frac{\overline{I^r J^s K^t}}{a^r \overline{J^s K^t} + b^s \overline{I^r K^t} + c^t \overline{I^r J^s}} \right) &= \lambda_R \left(\frac{R}{a^r \overline{J^s K^t} + b^s \overline{I^r K^t} + c^t \overline{I^r J^s}} \right) - \lambda_R \left(\frac{R}{\overline{I^r J^s K^t}} \right) \\ &= \lambda_R \left(\frac{R}{(a^r, b^s, c^t)} \right) + \lambda_R \left(\frac{(a^r, b^s, c^t)}{a^r \overline{J^s K^t} + b^s \overline{I^r K^t} + c^t \overline{I^r J^s}} \right) - \lambda_R \left(\frac{R}{\overline{I^r J^s K^t}} \right) \\ &= rst \lambda_R \left(\frac{R}{(a, b, c)} \right) + \left[\lambda_R \left(\frac{R}{\overline{I^r J^s}} \right) + \lambda_R \left(\frac{R}{\overline{I^r K^t}} \right) + \lambda_R \left(\frac{R}{\overline{J^s K^t}} \right) \right] \\ &- \left[\lambda_R \left(\frac{R}{\overline{I^r}} \right) + \lambda_R \left(\frac{R}{\overline{J^s}} \right) + \lambda_R \left(\frac{R}{\overline{K^t}} \right) \right] - \lambda_R \left(\frac{R}{\overline{I^r J^s K^t}} \right), \text{ by Proposition 4.3.} \end{aligned}$$

By [12, Theorem 2.4], $\bar{e}_{(1,1,1)} = \lambda_R \left(\frac{R}{(a, b, c)} \right)$ and

$\bar{e}_{(3,0)}(I, J) = \bar{e}_{(3,0)}(I, K) =$ $\bar{e}_{(3,0,0)} = e(I)$	$\bar{e}_{(0,3)}(I, J) = \bar{e}_{(0,3)}(J, K) =$ $\bar{e}_{(0,3,0)} = e(J)$	$\bar{e}_{(0,3)}(I, K) = \bar{e}_{(0,3)}(J, K) =$ $\bar{e}_{(0,0,3)} = e(K)$
$\bar{e}_{(2,1,0)} = \bar{e}_{(2,1)}(I, J)$	$\bar{e}_{(0,2,1)} = \bar{e}_{(2,1)}(J, K)$	$\bar{e}_{(2,0,1)} = \bar{e}_{(2,1)}(I, K)$
$\bar{e}_{(1,2,0)} = \bar{e}_{(1,2)}(I, J)$	$\bar{e}_{(0,1,2)} = \bar{e}_{(1,2)}(J, K)$	$\bar{e}_{(1,0,2)} = \bar{e}_{(1,2)}(I, K)$

Therefore, using Lemma 4.4, for $r, s, t \gg 0$, we have

$$\begin{aligned} \lambda_R \left(\frac{\overline{I^r J^s K^t}}{a^r \overline{J^s K^t} + b^s \overline{I^r K^t} + c^t \overline{I^r J^s}} \right) &= r[\bar{e}_{(1,0)}(I, J) + \bar{e}_{(1,0)}(I, K) - \bar{e}_{(1,0,0)} - \bar{e}_2(I)] \\ &+ s[\bar{e}_{(0,1)}(I, J) + \bar{e}_{(1,0)}(J, K) - \bar{e}_{(0,1,0)} - \bar{e}_2(J)] \\ &+ t[\bar{e}_{(0,1)}(I, K) + \bar{e}_{(0,1)}(J, K) - \bar{e}_{(0,0,1)} - \bar{e}_2(K)] \\ &+ \bar{e}_3(IJK) - [\bar{e}_3(IJ) + \bar{e}_3(IK) + \bar{e}_3(JK)] + \bar{e}_3(I) + \bar{e}_3(J) + \bar{e}_3(K). \end{aligned}$$

□

5. CHARACTERIZATION OF JOINT REDUCTION NUMBER ZERO IN TERMS OF LOCAL COHOMOLOGY MODULES

Let $H_K^i(M)$ denotes the local cohomology module of an R -module M with support in an ideal K of R . Let $(R, \mathfrak{m})$ be an analytically unramified local ring of dimension 3 and I, J, K be $\mathfrak{m}$ -primary ideals in R . Let $a \in I, b \in J$ and $c \in K$. Let $\overline{\mathcal{R}'}$ denote the extended Rees algebra of the filtration $\mathcal{F} = \{\overline{I^r J^s K^t}\}$. Consider the Koszul complex on $((at_1)^k, (bt_2)^k, (ct_3)^k)$ in $\overline{\mathcal{R}'}$:

$$F^{k\cdot} : 0 \longrightarrow \overline{\mathcal{R}'} \longrightarrow \bigoplus_{i=1}^3 \overline{\mathcal{R}'}(k\mathbf{e}_i) \xrightarrow{\alpha_k} \bigoplus_{1 \leq i < j \leq 3} \overline{\mathcal{R}'}(k(\mathbf{e}_i + \mathbf{e}_j)) \xrightarrow{\beta_k} \overline{\mathcal{R}'}(k(\mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3)) \longrightarrow 0.$$

The twists are given so that maps are degree zero maps. Then

$$H_{(at_1, bt_2, ct_3)}^i(\overline{\mathcal{R}'}) = \lim_{\xrightarrow{k}} H^i(F^{k\cdot}) \quad (22)$$

for all i by [1, Theorem 5.2.9]. The purpose of this section is to find the length of the component at origin of the third local cohomology module $H_{(at_1, bt_2, ct_3)}^3(\overline{\mathcal{R}'})$ in terms of normal Hilbert coefficients and joint reduction (a, b, c) . This leads to a satisfactory analogue of Rees's Theorem for 3-dimensional analytically unramified local rings.

Theorem 5.1. *Let $(R, \mathfrak{m})$ be an analytically unramified Cohen-Macaulay local ring of dimension 3 with infinite residue field and I, J, K be $\mathfrak{m}$ -primary ideals in R . Let (a, b, c) be a good joint reduction of $\{\overline{I^r J^s K^t}\}$. Then*

(1)

$$[H_{(at_1, bt_2, ct_3)}^3(\overline{\mathcal{R}'})]_{(0,0,0)} \cong \lim_{\xrightarrow{k}} \frac{\overline{I^k J^k K^k}}{a^k \overline{J^k K^k} + b^k \overline{I^k K^k} + c^k \overline{I^k J^k}}.$$

(2) For the directed system involved in the above direct limit, the map μ_k

$$\frac{\overline{I^k J^k K^k}}{a^k \overline{J^k K^k} + b^k \overline{I^k K^k} + c^k \overline{I^k J^k}} \xrightarrow{\mu_k} \frac{\overline{I^{k+1} J^{k+1} K^{k+1}}}{a^{k+1} \overline{J^{k+1} K^{k+1}} + b^{k+1} \overline{I^{k+1} K^{k+1}} + c^{k+1} \overline{I^{k+1} J^{k+1}}}$$

given by $\mu_k(x') = (xabc)'$, where $x \in \overline{I^k J^k K^k}$ and $'$ denotes the image of an element in respect quotients, is injective for all $k > 0$.

Proof. (1) The local cohomology module $H_{(at_1, bt_2, ct_3)}^3(\overline{\mathcal{R}'})$ has a $\mathbb{Z}^3$ -grading inherited from the $\mathbb{Z}^3$ -grading of $\overline{\mathcal{R}'}$. Therefore by the equation (22),

$$[H_{(at_1, bt_2, ct_3)}^3(\overline{\mathcal{R}'})]_{(0,0,0)} = \lim_{\rightarrow k} \frac{\overline{I^k J^k K^k}}{(\text{image } \beta_k)_{(0,0,0)}}.$$

Since $(\text{image } \beta_k)_{(0,0,0)} \simeq [a^k \overline{J^k K^k} + b^k \overline{I^k K^k} + c^k \overline{I^k J^k}]$,

$$[H_{(at_1, bt_2, ct_3)}^3(\overline{\mathcal{R}'})]_{(0,0,0)} \cong \lim_{\rightarrow k} \frac{\overline{I^k J^k K^k}}{a^k \overline{J^k K^k} + b^k \overline{I^k K^k} + c^k \overline{I^k J^k}}.$$

(2) Let $x \in \overline{I^k J^k K^k}$ be such that $\mu_k(x') = 0$. Then $xabc = a^{k+1}u + b^{k+1}v + c^{k+1}w$ for some $u \in \overline{J^{k+1} K^{k+1}}$, $v \in \overline{I^{k+1} K^{k+1}}$ and $w \in \overline{I^{k+1} J^{k+1}}$. Hence $u \in (b, c) \cap \overline{J^{k+1} K^{k+1}} = b \overline{J^k K^{k+1}} + c \overline{J^{k+1} K^k}$. Thus $u = bu_1 + cu_2$ for some $u_1 \in \overline{J^k K^{k+1}}$ and $u_2 \in \overline{J^{k+1} K^k}$. Similarly, $v = av_1 + cv_2$, $w = aw_1 + bw_2$ for some $v_1 \in \overline{I^k K^{k+1}}$, $v_2 \in \overline{I^{k+1} K^k}$, $w_1 \in \overline{I^k J^{k+1}}$ and $w_2 \in \overline{I^{k+1} J^k}$. Thus,

$$xabc = ac(a^k u_2 + c^k w_1) + bc(b^k v_2 + c^k w_2) + ab(a^k u_1 + b^k v_1).$$

Hence $a^k u_1 + b^k v_1 \in (c) \cap \overline{I^k J^k K^{k+1}} = c \overline{I^k J^k K^k}$. Let $a^k u_1 + b^k v_1 = cp$ for some $p \in \overline{I^k J^k K^k}$. Then $p \in (a^k, b^k) \cap \overline{I^k J^k K^k} = a^k \overline{J^k K^k} + b^k \overline{I^k K^k}$. Similarly, $a^k u_2 + c^k w_1 = bq$ for some $q \in a^k \overline{J^k K^k} + c^k \overline{I^k J^k}$ and $b^k v_2 + c^k w_2 = ar$ for some $r \in b^k \overline{I^k K^k} + c^k \overline{I^k J^k}$. Hence $x = p + q + r \in a^k \overline{J^k K^k} + b^k \overline{I^k K^k} + c^k \overline{I^k J^k}$. Thus $x' = 0$ and therefore μ_k is injective. $\square$

Lemma 5.2. Let $(R, \mathfrak{m})$ be a Noetherian local ring. Let $\{(M_i, \psi_{i,j})\}_{i \geq 0}$ be a directed system of R -modules and $M = \varinjlim_k M_k$.

(1) If M is a finitely generated R -module then the natural map $\psi_k : M_k \rightarrow M$ is surjective for all large k .

(2) If $\psi_{k,k+1} : M_k \rightarrow M_{k+1}$ is injective for $k \gg 0$ then $\lambda_k : M_k \rightarrow M$ is injective for large k .

Proof. (1) Let $\lambda_i : M_i \rightarrow \bigoplus_{i \geq 0} M_i$ be the i th injection. We know that $M = \frac{\bigoplus_k M_k}{N}$, where N is the R -submodule of $\bigoplus_k M_k$ generated by elements $\lambda_j \psi_{i,j}(m_i) - \lambda_i(m_i)$ for $m_i \in M_i$ and $i \leq j$. Let $\psi_k : M_k \rightarrow M$ be the natural map defined as $\psi_k(m_k) = \lambda_k(m_k) + N$. Suppose that M is generated by $\psi_{k_1}(x_{k_1}), \psi_{k_2}(x_{k_2}), \dots, \psi_{k_g}(x_{k_g})$ for some $x_{k_i} \in M_{k_i}$. Let $k_0 = \max\{k_1, k_2, \dots, k_g\}$. Since $\psi_{k_i} = \psi_k \psi_{k_i,k}$ for $k \geq k_0$, M is also generated by $\psi_k(\psi_{k_i,k}(x_{k_i}))$ for $i = 1, 2, \dots, g$. Hence $\psi_k : M_k \rightarrow M$ is surjective for $k \geq k_0$.

(2) Suppose $\psi_k(x) = 0$ for some $x \in M_k$. Then $\lambda_k(x) + N = 0$. By [14, Lemma 5.30], $\psi_{k,j}(x) = 0$ for some $j \geq k$. Hence $x = 0$ if $k \gg 0$. $\square$

Theorem 5.3. *Let $(R, \mathfrak{m})$ be an analytically unramified Cohen-Macaulay local ring of dimension 3 with infinite residue field and I, J, K be $\mathfrak{m}$ -primary ideals in R . Assume that a good complete reduction of the filtration $\{\overline{I^r J^s K^t}\}$ exists. Let (a, b, c) be any good joint reduction of the filtration $\{\overline{I^r J^s K^t}\}$. Then*

- (1) $\lambda_R([H_{(at_1, bt_2, ct_3)}^3(\overline{\mathcal{R}'})]_{(0,0,0)}) < \infty$.
- (2) $\lambda_R([H_{(at_1, bt_2, ct_3)}^3(\overline{\mathcal{R}'})]_{(0,0,0)}) = \lambda_R\left(\frac{\overline{I^r J^s K^t}}{a^r \overline{J^s K^t} + b^s \overline{I^r K^t} + c^t \overline{I^r J^s}}\right)$ for large r, s, t .
- (3) $\lambda_R([H_{(at_1, bt_2, ct_3)}^3(\overline{\mathcal{R}'})]_{(0,0,0)}) = \bar{e}_3(IJK) - [\bar{e}_3(IJ) + \bar{e}_3(IK) + \bar{e}_3(JK)] + \bar{e}_3(I) + \bar{e}_3(J) + \bar{e}_3(K)$.

In particular, $\lambda_R([H_{(at_1, bt_2, ct_3)}^3(\overline{\mathcal{R}'})]_{(0,0,0)})$ is independent of good joint reduction (a, b, c) of $\{\overline{I^r J^s K^t}\}$.

Proof. (1) By Theorem 5.1(1),

$$[H_{(at_1, bt_2, ct_3)}^3(\overline{\mathcal{R}'})]_{(0,0,0)} \cong \lim_{\rightarrow} \frac{\overline{I^k J^k K^k}}{a^k \overline{J^k K^k} + b^k \overline{I^k K^k} + c^k \overline{I^k J^k}}.$$

Let $S_{(a,b,c)}(r, s, t) = \lambda_R\left(\frac{\overline{I^r J^s K^t}}{a^r \overline{J^s K^t} + b^s \overline{I^r K^t} + c^t \overline{I^r J^s}}\right)$. First we show that $S_{(a,b,c)}(r, s, t)$ is a constant for $r, s, t \gg 0$. Let (x_{ij}) be a good complete reduction of $\{\overline{I^r J^s K^t}\}$. Let $x = x_{11}, y = x_{22}$ and $z = x_{33}$. Then, by Remark 2.6(2), (x, y, z) is a good joint reduction of $\{\overline{I^r J^s K^t}\}$. By Theorem 2.8, $\lambda_R([H_{(xt_1, yt_2, zt_3)}^3(\overline{\mathcal{R}'})]_{(0,0,0)}) < \infty$. Thus $[H_{(xt_1, yt_2, zt_3)}^3(\overline{\mathcal{R}'})]_{(0,0,0)}$ is a finitely generated R -module. Therefore, by Theorem 5.1 and Lemma 5.2, for $k \gg 0$,

$$[H_{(xt_1, yt_2, zt_3)}^3(\overline{\mathcal{R}'})]_{(0,0,0)} \cong \frac{\overline{I^k J^k K^k}}{x^k \overline{J^k K^k} + y^k \overline{I^k K^k} + z^k \overline{I^k J^k}}$$

Hence

$$\lambda_R([H_{(xt_1, yt_2, zt_3)}^3(\overline{\mathcal{R}'})]_{(0,0,0)}) = S_{(x,y,z)}(k, k, k). \quad (23)$$

By Theorem 4.5, $S_{(x,y,z)}(r, s, t)$ is a polynomial function in r, s, t of total degree atmost 1. Fixing s_0, t_0 large, $S_{(x,y,z)}(r, s_0, t_0)$ is a polynomial function in r of degree atmost 1. Hence coefficient of r in the polynomial associated with $S_{(x,y,z)}(r, s_0, t_0)$ is nonnegative. Thus the coefficient of r in the polynomial associated with $S_{(x,y,z)}(r, s, t)$ is nonnegative. Similarly, coefficients of s and t in the polynomial associated with $S_{(x,y,z)}(r, s, t)$ are nonnegative. By equation (23), $S_{(x,y,z)}(k, k, k)$ is a constant for large k . Hence the sum of coefficients of r, s, t in the polynomial associated with $S_{(x,y,z)}(r, s, t)$ is zero. Hence each coefficient of r, s, t in the polynomial associated with $S_{(x,y,z)}(r, s, t)$ is zero. Thus, by Theorem 4.5, for $r, s, t \gg 0$,

$$S_{(a,b,c)}(r, s, t) = \bar{e}_3(IJK) - [\bar{e}_3(IJ) + \bar{e}_3(IK) + \bar{e}_3(JK)] + \bar{e}_3(I) + \bar{e}_3(J) + \bar{e}_3(K) \quad (24)$$

for any good joint reduction (a, b, c) of $\{\overline{I^r J^s K^t}\}$. Let

$$\mu_k : \frac{\overline{I^k J^k K^k}}{a^k \overline{J^k K^k} + b^k \overline{I^k K^k} + c^k \overline{I^k J^k}} \longrightarrow \frac{\overline{I^{k+1} J^{k+1} K^{k+1}}}{a^{k+1} \overline{J^{k+1} K^{k+1}} + b^{k+1} \overline{I^{k+1} K^{k+1}} + c^{k+1} \overline{I^{k+1} J^{k+1}}}$$

be the map involved in the direct limit. By Theorem 5.1(2), μ_k is injective for $k > 0$.

By equation (24), $S_{(a,b,c)}(k, k, k)$ is a constant for $k \gg 0$. Hence μ_k is an isomorphism for $k \gg 0$. Therefore for $k \gg 0$,

$$[H_{(at_1, bt_2, ct_3)}^3(\overline{\mathcal{R}'})]_{(0,0,0)} \cong \frac{\overline{I^k J^k K^k}}{a^k \overline{J^k K^k} + b^k \overline{I^k K^k} + c^k \overline{I^k J^k}} \quad (25)$$

and hence $[H_{(at_1, bt_2, ct_3)}^3(\overline{\mathcal{R}'})]_{(0,0,0)}$ has finite length.

- (2) Follows from equations (24) and (25).
- (3) Follows from equations (24) and (25).

□

Theorem 5.4. *Let $(R, \mathfrak{m})$ be an analytically unramified Cohen-Macaulay local ring of dimension 3 and I, J, K be $\mathfrak{m}$ -primary ideals in R . Assume that a good complete reduction of the filtration $\{\overline{I^r J^s K^t}\}$ exists. The following statements are equivalent:*

- (1) $[H_{(at_1, bt_2, ct_3)}^3(\overline{\mathcal{R}'})]_{(0,0,0)} = 0$ for some good joint reduction (a, b, c) of the filtration $\{\overline{I^r J^s K^t}\}$,
- (2) $[H_{(at_1, bt_2, ct_3)}^3(\overline{\mathcal{R}'})]_{(0,0,0)} = 0$ for any good joint reduction (a, b, c) of the filtration $\{\overline{I^r J^s K^t}\}$,
- (3) the normal joint reduction number of I, J, K is zero with respect to any good joint reduction (a, b, c) of the filtration $\{\overline{I^r J^s K^t}\}$,
- (4) the normal joint reduction number of I, J, K is zero with respect to some good joint reduction (a, b, c) of the filtration $\{\overline{I^r J^s K^t}\}$,

$$(5) \quad \bar{e}_3(IJK) - [\bar{e}_3(IJ) + \bar{e}_3(IK) + \bar{e}_3(JK)] + \bar{e}_3(I) + \bar{e}_3(J) + \bar{e}_3(K) = 0.$$

Proof. (1) $\Rightarrow$ (2) : The result follows by the Theorem 5.3(3).

(2) $\Rightarrow$ (3) : By Theorem 5.3(2), for $r, s, t \gg 0$,

$$\overline{I^r J^s K^t} = a^r \overline{J^s K^t} + b^s \overline{I^r K^t} + c^t \overline{I^r J^s},$$

say for $r, s, t \geq N \geq 1$. Suppose $N \geq 2$. We prove that for $s, t \geq N$

$$\overline{I^{N-1} J^s K^t} = a^{N-1} \overline{J^s K^t} + b^s \overline{I^{N-1} K^t} + c^t \overline{I^{N-1} J^s}.$$

Let $z \in \overline{I^{N-1} J^s K^t}$. Then $az \in \overline{I^N J^s K^t}$. Let $az = a^N u + b^s v + c^t w$ for some $u \in \overline{J^s K^t}, v \in \overline{I^N K^t}, w \in \overline{I^N J^s}$. Then $z - a^{N-1} u \in (b^s, c^t) \cap \overline{I^{N-1} J^s K^t} = b^s \overline{I^{N-1} K^t} + c^t \overline{I^{N-1} J^s}$, by Lemma 4.1. Hence $z \in a^{N-1} \overline{J^s K^t} + b^s \overline{I^{N-1} K^t} + c^t \overline{I^{N-1} J^s}$.

Similar argument shows that

$$\begin{aligned} \overline{I^r J^{N-1} K^t} &= a^r \overline{J^{N-1} K^t} + b^{N-1} \overline{I^r K^t} + c^t \overline{I^r J^{N-1}} \text{ for } r, t \geq N \\ \overline{I^r J^s K^{N-1}} &= a^r \overline{J^s K^{N-1}} + b^s \overline{I^r K^{N-1}} + c^{N-1} \overline{I^r J^s} \text{ for } r, s \geq N. \end{aligned}$$

Continuing as above, we get that for all $r, s, t \geq 1$,

$$\overline{I^r J^s K^t} = a^r \overline{J^s K^t} + b^s \overline{I^r K^t} + c^t \overline{I^r J^s}.$$

Hence $\overline{I^r J^s K^t} \subseteq a \overline{I^{r-1} J^s K^t} + b \overline{I^r J^{s-1} K^t} + c \overline{I^r J^s K^{t-1}}$ for all $r, s, t \geq 1$. Thus the normal joint reduction number of I, J, K is zero with respect to (a, b, c) .

(3) $\Rightarrow$ (4) : This is clear.

(4) $\Rightarrow$ (5) : First we prove that for $r, s, t > 0$,

$$\overline{I^r J^s K^t} = a^r \overline{J^s K^t} + b \overline{I^r J^{s-1} K^t} + c \overline{I^r J^s K^{t-1}}. \quad (26)$$

Induct on r . Since the normal joint reduction number of I, J, K is zero with respect to (a, b, c) , the result is true for $r = 1$. Let $r > 1$ and $x \in \overline{I^r J^s K^t}$. Therefore $x = au + bv + cw$ for some $u \in \overline{I^{r-1} J^s K^t}, v \in \overline{I^r J^{s-1} K^t}, w \in \overline{I^r J^s K^{t-1}}$. By induction, $u = a^{r-1} u_1 + b u_2 + c u_3$ for some $u_1 \in \overline{J^s K^t}, u_2 \in \overline{I^{r-1} J^{s-1} K^t}, u_3 \in \overline{I^{r-1} J^s K^{t-1}}$. Hence $x \in a^r \overline{J^s K^t} + b \overline{I^r J^{s-1} K^t} + c \overline{I^r J^s K^{t-1}}$. Similarly, for $r, s, t > 0$,

$$\overline{I^r J^s K^t} = a \overline{I^{r-1} J^s K^t} + b^s \overline{I^r K^t} + c \overline{I^r J^s K^{t-1}} \text{ and} \quad (27)$$

$$\overline{I^r J^s K^t} = a \overline{I^{r-1} J^s K^t} + b \overline{I^r J^{s-1} K^t} + c^t \overline{I^r J^s}. \quad (28)$$

We use induction on $r + s + t$ to prove that for $r, s, t > 0$,

$$\overline{I^r J^s K^t} = a^r \overline{J^s K^t} + b^s \overline{I^r K^t} + c^t \overline{I^r J^s}.$$

Suppose $r + s + t = 3$. Then $r = s = t = 1$. Since the joint reduction number is zero, result is true in this case. Let $r + s + t > 3$. Without loss of generality we may assume that $s > 1$. If $r = t = 1$, then result follows from the equation (27). Therefore we may also assume that $t > 1$. Let $x \in \overline{I^r J^s K^t}$. Then, by the equation (26), $x = a^r u + bv + cw$ for some $u \in \overline{J^s K^t}$, $v \in \overline{I^r J^{s-1} K^t}$ and $w \in \overline{I^r J^s K^{t-1}}$. By induction, $v = a^r v_1 + b^{s-1} v_2 + c^t v_3$ and $w = a^r w_1 + b^s w_2 + c^{t-1} w_3$ for some $v_1 \in \overline{J^{s-1} K^t}$, $v_2 \in \overline{I^r K^t}$, $v_3 \in \overline{I^r J^{s-1}}$, $w_1 \in \overline{J^s K^{t-1}}$, $w_2 \in \overline{I^r K^{t-1}}$, $w_3 \in \overline{I^r J^s}$. Thus $x \in a^r \overline{J^s K^t} + b^s \overline{I^r K^t} + c^t \overline{I^r J^s}$.

Therefore, by Theorem 4.5, result follows.

(5) $\Rightarrow$ (1) : Follows from Theorem 5.3(3). $\square$

Theorem 5.5. *Let $(R, \mathfrak{m})$ be an analytically unramified Cohen-Macaulay local ring of dimension 3 and I, J, K be $\mathfrak{m}$ -primary ideals in R . Let (a, b, c) be a good joint reduction of the filtration $\{\overline{I^r J^s K^t}\}$. Assume that $\lambda_R(R/\overline{I^r J^s K^t}) = \overline{P}_{I,J,K}(r, s, t)$ for $r + s + t > 0$. Then the normal joint reduction number of I, J, K is zero with respect to (a, b, c) if and only of $\overline{e}_3(IJK) = 0$.*

Proof. Since $\lambda_R(R/\overline{I^r J^s K^t}) = \overline{P}_{I,J,K}(r, s, t)$ for $r + s + t > 0$, taking $s = t = 0$ and $r > 0$, we get

$$\overline{e}_{(1,0,0)} = \overline{e}_2(I), \overline{e}_3(IJK) = \overline{e}_3(I).$$

Taking $t = 0$ and $r, s \gg 0$, we get $\overline{e}_{(1,0,0)} = \overline{e}_{(1,0)}(I, J)$, $\overline{e}_{(0,1,0)} = \overline{e}_{(0,1)}(I, J)$ and $\overline{e}_3(IJK) = \overline{e}_3(IJ)$.

Since $\lambda_R(R/\overline{I^r J^s K^t}) = \overline{P}_{I,J,K}(r, s, t)$ for $r + s + t > 0$, arguing similarly and using Theorem 4.5, we get that for $r, s, t > 0$,

$$\lambda_R \left(\frac{\overline{I^r J^s K^t}}{a^r \overline{J^s K^t} + b^s \overline{I^r K^t} + c^t \overline{I^r J^s}} \right) = \overline{e}_3(IJK).$$

Thus $\overline{e}_3(IJK) = 0$ if and only if for $r, s, t > 0$, $\overline{I^r J^s K^t} = a^r \overline{J^s K^t} + b^s \overline{I^r K^t} + c^t \overline{I^r J^s}$. Hence the normal joint reduction number of I, J, K is zero with respect to (a, b, c) if and only of $\overline{e}_3(IJK) = 0$. $\square$

6. SOME APPLICATIONS

In this section we discuss a few applications of Theorem 5.4. Let $(R, \mathfrak{m})$ be an analytically unramified local ring of dimension d and I be an $\mathfrak{m}$ -primary ideal in R . An ideal $K \subseteq I$ is said to

be a *reduction* of the filtration $\{\overline{I^n}\}$ if $K\overline{I^n} = \overline{I^{n+1}}$ for all large n . A *minimal reduction* of $\{\overline{I^n}\}$ is a reduction of $\{\overline{I^n}\}$ minimal with respect to inclusion. For a minimal reduction K of $\{\overline{I^n}\}$, we set

$$\bar{\tau}_K(I) = \sup\{n \in \mathbb{Z} \mid \overline{I^n} \neq K\overline{I^{n-1}}\}.$$

The *reduction number* $\bar{\tau}(I)$ of $\{\overline{I^n}\}$ is defined to be the least $\bar{\tau}_K(I)$ over all possible minimal reductions K of $\{\overline{I^n}\}$.

Theorem 6.1. [9, Corollary 3.17] *Let $(R, \mathfrak{m})$ be an analytically unramified Cohen-Macaulay local ring of dimension $d \geq 3$ with infinite residue field and I be an $\mathfrak{m}$ -primary ideal in R . Let $\overline{G}(I) = \bigoplus_{n \geq 0} \overline{I^n}/\overline{I^{n+1}}$ be the associated graded algebra of $\{\overline{I^n}\}$. Suppose $\text{depth } \overline{G}(I) \geq d-1$. Then $\bar{\tau}(I) \leq 2$ if and only if $\bar{e}_3(I) = 0$.*

Proof. Let $\bar{\tau}(I) \leq 2$. Then $\bar{e}_3(I) = 0$ by [9, Corollary 3.17]. To prove the converse we use induction on d . Let $d = 3$. Since $\text{depth } \overline{G}(I) \geq 2$, there exists a minimal reduction $J = (a, b, c)$ of $\{\overline{I^n}\}$ such that a^*, b^*, c^* are nonzerodivisors in $\overline{G}(I)$ and $(a^*, b^*), (b^*, c^*), (a^*, c^*)$ are $\overline{G}(I)$ -regular sequences. Hence (x_{ij}) , where $x_{i1} = a, x_{i2} = b$ and $x_{i3} = c$, for all $i = 1, 2, 3$, is a good complete reduction of $\{\overline{I^r I^s I^t}\}$ and (a, b, c) is a good joint reduction of $\{\overline{I^r I^s I^t}\}$. Since $\bar{e}_3(I) = \bar{e}_3(I^2) = \bar{e}_3(I^3)$, by Theorem 5.4, $\bar{e}_3(I) = 0$ if and only if $\bar{\tau}_J(I) \leq 2$. By [9, Corollary 3.8], $\bar{\tau}(I)$ is independent of the minimal reduction chosen. Hence $\bar{e}_3(I) = 0$ if and only if $\bar{\tau}(I) \leq 2$.

Suppose $d > 3$. Let $I = (a_1, \dots, a_l)$ for some $l \geq 1$. For indeterminates $X_1, \dots, X_l$, set $S = R[X_1, \dots, X_l]_{\mathfrak{m}[X_1, \dots, X_l]}$, $z = a_1 X_1 + \dots + a_l X_l$ and $C = S/zC$. Then, by [6, Theorem 1] and [6, Corollary 8], C is an analytically unramified Cohen-Macaulay local ring of dimension $d-1$ and $\bar{e}_3(I) = \bar{e}_3(IC)$. Therefore then $\bar{\tau}(IC) \leq 2$, by induction. Hence, by [6, Proposition 17], $\bar{\tau}(I) \leq 2$. $\square$

Theorem 6.2. *Let $(R, \mathfrak{m})$ be a Cohen-Macaulay local ring of dimension d and $\mathbf{I} = (I_1, I_2, I_3)$ be $\mathfrak{m}$ -primary ideals in R . Let $\mathcal{F}$ be an $\mathbf{I}$ -admissible filtration. Suppose that $\mathcal{R}(\mathcal{F})$ is Cohen-Macaulay. Then $P_{\mathcal{F}}(\mathbf{n}) = H_{\mathcal{F}}(\mathbf{n})$ for all $\mathbf{n} \geq \mathbf{0}$.*

Proof. Since $a^j(\mathcal{R}(\mathcal{F})) = -1$, by Lemma 3.7, for each $j = 1, 2, 3$

$$[H_{\mathcal{M}}^i(\mathcal{R}(\mathcal{F}))]_{\mathbf{n}} = 0 \text{ for all } n_j \geq 0 \text{ and all } i \geq 0.$$

Therefore, by Lemma 3.5, $[H_{\mathcal{R}_{++}}^i(\mathcal{R}(\mathcal{F}))]_{\mathbf{n}} = 0$ for all $\mathbf{n} \geq \mathbf{0}$ and all $i \geq 0$. Thus, by Lemma 3.8, $[H_{\mathcal{R}_{++}}^i(\mathcal{R}'(\mathcal{F}))]_{\mathbf{n}} = 0$ for all $\mathbf{n} \geq \mathbf{0}$ and all $i \geq 0$. Hence, by difference formula [7, Theorem 5.1], for

all $\mathbf{n} \geq \mathbf{0}$, we have

$$P_{\mathcal{F}}(\mathbf{n}) - H_{\mathcal{F}}(\mathbf{n}) = \sum_{i=0}^d (-1)^i \lambda_R([H_{\mathcal{R}^{++}}^i(\mathcal{R}'(\mathcal{F}))]_{\mathbf{n}}) = 0$$

Hence $P_{\mathcal{F}}(\mathbf{n}) = H_{\mathcal{F}}(\mathbf{n})$ for all $\mathbf{n} \geq \mathbf{0}$. $\square$

Theorem 6.3. *Let $R = k[x, y, z]$ be polynomial ring in 3 variables x, y, z . Let $\mathfrak{m} = (x, y, z)$ and I, J, K be an $\mathfrak{m}$ -primary monomial ideal in R . Let $S = R_{\mathfrak{m}}$. Then the normal joint reduction number of IS, JS, KS is zero with respect to every good joint reduction of $\{\overline{I^r J^s K^t S}\}$.*

Proof. Note that S is an analytically unramified Cohen-Macaulay local ring of dimension 3 and IS, JS, KS are $\mathfrak{m}S$ -primary ideals in S . By [2, Theorem 6.3.5], $\overline{\mathcal{R}}(I, J, K)$ is Cohen-Macaulay. Since $\overline{\mathfrak{a}S} = \overline{\mathfrak{a}}S$ for every ideal $\mathfrak{a}$ in R , we have

$$\overline{\mathcal{R}}(IS, JS, KS) = \bigoplus_{r,s,t \geq 0} \overline{I^r J^s K^t S t_1^r t_2^s t_3^t} = \bigoplus_{r,s,t \geq 0} \overline{I^r J^s K^t S} t_1^r t_2^s t_3^t = \overline{\mathcal{R}}(I, J, K)S.$$

Hence $\overline{\mathcal{R}}(IS, JS, KS)$ is Cohen-Macaulay. Thus, by Theorem 3.9, there exists a good complete reduction $(x_{ij})_{1 \leq i, j \leq 3}$ of the filtration $\{\overline{I^r J^s K^t S}\}$. By Theorem 6.2, $\overline{P}_{IS, JS, KS}(r, s, t) = \overline{H}_{IS, JS, KS}(r, s, t)$ for all $r, s, t \geq 0$. Hence $\overline{P}_{IS, JS, KS}(0, 0, 0) = \overline{e}_3(IJKS) = 0$. Therefore, by Theorem 5.5, the normal joint reduction number of IS, JS, KS is zero with respect to every good joint reduction (a, b, c) of $\{\overline{I^r J^s K^t S}\}$. $\square$

Theorem 6.4. *Let $R = k[x, y, z]$, $\mathfrak{m} = (x, y, z)$ and I be an $\mathfrak{m}$ -primary monomial ideal in R . Suppose I and I^2 are complete. Then I^n is complete for all $n \geq 1$.*

Proof. Let $S = R_{\mathfrak{m}}$. Then, by Theorem 6.3, the normal joint reduction number of IS, IS, IS is zero with respect to every good joint reduction $J = (a, b, c)$ of $\{\overline{I^r I^s I^t S}\}$. Hence $\overline{I^n S} = J \overline{I^{n-1} S}$ for all $n \geq 3$. Since I and I^2 are complete, IS and $I^2 S$ are complete. Thus $I^n S$ is complete for all $n \geq 1$. Hence I^n is complete for all $n \geq 1$. $\square$

Theorem 6.5. *Let $R = k[x, y, z]$, $\mathfrak{m} = (x, y, z)$ and I, J be $\mathfrak{m}$ -primary monomial ideals in R . Suppose that $I^r J^s$ is complete for all $r, s \geq 0$ such that $r + s \leq 2$. Then $I^r J^s$ is complete for all $r, s \geq 0$.*

Proof. Let $S = R_{\mathfrak{m}}$. By Theorem 6.3, the normal joint reduction number of IS, IS, JS and IS, JS, JS is zero with respect to every good joint reduction of $\{\overline{I^r I^s J^t S}\}$ and $\{\overline{I^r J^s J^t S}\}$ respectively. Let (a_1, b_1, c_1) be a good joint reduction of the filtration $\{\overline{I^r I^s J^t S}\}$ and (a_2, b_2, c_2) be a good joint reduction of the filtration $\{\overline{I^r J^s J^t S}\}$. Then

$$\overline{I^r J^s S} = (a_1, b_1) \overline{I^{r-1} J^s S} + c_1 \overline{I^r J^{s-1} S} \text{ for all } (r, s) \geq (2, 1) \text{ and} \quad (29)$$

$$\overline{I^r J^s S} = (a_2) \overline{I^{r-1} J^s S} + (b_2, c_2) \overline{I^r J^{s-1} S} \text{ for all } (r, s) \geq (1, 2). \quad (30)$$

It suffices to show that $I^r J^s S$ is complete for all $r, s \geq 0$. By Theorem 6.4, $I^n S$ and $J^n S$ are complete for all $n \geq 1$. We use induction on $r + s$ to show that $I^r J^s S$ is complete for all $r, s \geq 0$. By assumption, result is true for $r + s \leq 2$. Suppose $r + s > 2$. We may assume that $r > 1$. Since $I^r S$ is complete for all $r \geq 1$, assertion is true for $s = 0$. Hence we may also assume that $s \geq 1$. By induction, $I^{r-1} J^s S$ and $I^r J^{s-1} S$ are complete. Therefore, by equation (29), $I^r J^s S$ is complete. $\square$

Theorem 6.6. *Let $R = k[x, y, z]$, $\mathfrak{m} = (x, y, z)$ and I, J, K be $\mathfrak{m}$ -primary monomial ideals in R . Suppose that $I^r J^s K^t$ is complete for all $r, s, t \geq 0$ such that $r + s + t \leq 2$. Then $I^r J^s K^t$ is complete for all $r, s, t \geq 0$.*

Proof. Let $S = R_{\mathfrak{m}}$. By Theorem 6.3, the normal joint reduction number of IS, JS, KS is zero with respect to every good joint reduction of the filtration $\{\overline{I^r J^s K^t S}\}$. Let (a, b, c) be a good joint reduction of the filtration $\{\overline{I^r J^s K^t S}\}$. Hence, for all $r, s, t \geq 1$,

$$\overline{I^r J^s K^t S} = a \overline{I^{r-1} J^s K^t S} + b \overline{I^r J^{s-1} K^t S} + c \overline{I^r J^s K^{t-1} S}. \quad (31)$$

It suffices to show that $I^r J^s K^t S$ is complete for all $r, s, t \geq 0$. We use induction on $r + s + t$ to show that $I^r J^s K^t S$ is complete for all $r, s, t \geq 0$. By assumption result is true for $r + s + t \leq 2$. Let $r + s + t \geq 3$. By Theorem 6.5, $I^r J^s S, I^r K^t S$ and $J^s K^t S$ are complete for all $r, s, t \geq 0$. Hence result is true if $r = 0$ or $s = 0$ or $t = 0$. Therefore, we may assume that $r, s, t \geq 1$. By induction, $I^{r-1} J^s K^t JS, I^r J^{s-1} K^t S$ and $I^r J^s K^{t-1} S$ are complete. Hence, by equation (31), $I^r J^s K^t S$ is complete. $\square$

REFERENCES

- [1] M. P. Brodmann and R. Y. Sharp, *Local cohomology : An algebraic introduction with geometric applications*, Second edition, Cambridge University Press, (2013). 6, 30
- [2] W. Bruns and J. Herzog, *Cohen-Macaulay Rings*, Revised Edition, Cambridge University Press, (1998). 37

- [3] C. D'Cruz and S. K. Masuti, *Local Cohomology of bigraded Rees algebras, Bhattacharya Coefficients and Joint Reductions*, [arXiv:1405.1550](#). [12](#)
- [4] M. Herrmann, E. Hyry and J. Ribbe, *On the Cohen-Macaulay and Gorenstein properties of multigraded Rees algebras*, Manuscripta Math. **79** (1993), 343-377. [21](#)
- [5] E. Hyry, *The diagonal subring and the Cohen-Macaulay property of a multigraded ring*, Trans. Amer. Math. Soc. **351** (1999), 2213-2232. [20](#), [21](#), [22](#)
- [6] S. Itoh, *Coefficients of normal Hilbert polynomials*, J. Algebra **150** (1992), 101-117. [36](#)
- [7] A. V. Jayanthan and J. K. Verma, *Grothendieck-Serre formula and bigraded Cohen-Macaulay Rees algebras*, J. Algebra **254** (2002), 1-20. [7](#), [8](#), [36](#)
- [8] D. Kirby and H. A. Mehran, *Hilbert functions and the Koszul complex*, J. London Math. Soc. (2) **24** (1981), 459-466. [24](#)
- [9] T. Marley, *Hilbert functions of ideals in Cohen-Macaulay rings*, Ph. D. Thesis, Purdue University, (1989). [36](#)
- [10] S. K. Masuti, *Normal Hilbert coefficients and bigraded Rees algebras*, Ph. D. Thesis, Indian Institute of Technology Bombay, (2013). [29](#)
- [11] S. K. Masuti and J. K. Verma, *Local cohomology of bigraded Rees algebras and normal Hilbert coefficients*, J. Pure Appl. Algebra **218** (2014), 904-918. [5](#)
- [12] D. Rees, *Generalizations of reductions and mixed multiplicities*, J. London Math. Soc. (2) **29** (1984), 397-414. [12](#), [28](#), [29](#)
- [13] D. Rees, *Hilbert functions and pseudo-rational local rings of dimension two*, J. London Math. Soc. (2) **24** (1981), 467-479. [1](#), [4](#), [28](#), [29](#)
- [14] J. Rotman, *An introduction to homological algebra*, Second edition, Springer, New York, (2009). [32](#)
- [15] M. A. Vitulli, *Some normal monomial ideals*, Contemp. Math. **324** (2003), 205-217. [2](#), [5](#)

INSTITUTE OF MATHEMATICAL SCIENCES, IV CROSS ROAD, CIT CAMPUS, TARAMANI, CHENNAI 600113, TAMIL NADU, INDIA.

E-mail address: shreedevikm@imsc.res.in

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY BOMBAY, POWAI, MUMBAI, INDIA - 400076

E-mail address: tputhen@math.iitb.ac.in

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY BOMBAY, POWAI, MUMBAI, INDIA - 400076

E-mail address: jkv@math.iitb.ac.in